

Generalized Fesenko reciprocity map

Kâzım İlhan İKEDA and Erol SERBEST

Abstract

In this paper, which is the natural continuation and generalization of [1, 2, 3] and [8], we extend the theory of Fesenko to infinite *APF*-Galois extensions L over a local field K , with finite residue-class field κ_K of $q = p^f$ elements, satisfying $\mu_p(K^{sep}) \subset K$ and $K \subset L \subset K_{\varphi^d}$ where the residue-class degree $[\kappa_L : \kappa_K] = d$. More precisely, for such extensions L/K , fixing a Lubin-Tate splitting φ over K , we construct a 1-cocycle,

$$\Phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \rightarrow K^\times / N_{L_0/K} L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0},$$

where $L_0 = L \cap K^{nr}$, and study its functorial and ramification-theoretic properties. The case $d = 1$ recovers the theory of Fesenko.

2000 Mathematics Subject Classification : Primary 11S37

Keywords : Local fields, higher-ramification theory, *APF*-extensions, Fontaine-Wintenberger field of norms, Fesenko reciprocity map, generalized Fesenko reciprocity map, non-abelian local class field theory.

Let K be a local field (that is, a complete discrete valuation field) with finite residue-class field κ_K of $q = p^f$ elements. Assume that $\mu_p(K^{sep}) \subset K$. Fix a Lubin-Tate splitting φ over K (cf. [10]). In [1, 2, 3], Fesenko introduced a very general non-abelian local reciprocity map

$$\Phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \rightarrow U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/K}$$

defined for any totally-ramified infinite *APF*-Galois extension L/K satisfying $K \subset L \subset K_\varphi$, which generalizes the previous non-abelian local class field theories of Koch-de Shalit [10] and Gurevich [7]. In [8], we have studied the basic functorial and ramification-theoretic properties of the reciprocity map of Fesenko.

In this paper, which is the natural continuation and generalization of [1, 2, 3] and [8], we extend the theory of Fesenko to infinite *APF*-Galois extensions L/K satisfying $K \subset L \subset K_{\varphi^d}$ where the residue-class degree $[\kappa_L : \kappa_K] = d$. More precisely, for such extensions L/K , we construct a 1-cocycle,

$$\Phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \rightarrow K^\times / N_{L_0/K} L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0},$$

where $L_0 = L \cap K^{nr}$, and study its functorial and ramification-theoretic properties. Note that, the case $d = 1$ recovers the theory of Fesenko.

The organization of this paper is as follows. In the first section, we briefly review the necessary background material from Fontaine-Wintenberger theory of fields of norms. In the second section, we introduce the generalized Fesenko reciprocity map $\Phi_{L/K}^{(\varphi)}$ of an extension L/K , which is an infinite *APF*-Galois

extension satisfying $K \subset L \subset K_{\varphi^d}$ where the residue-class degree $[\kappa_L : \kappa_K] = d$, and study its functorial and ramification-theoretic properties.

The material and results of this paper play a fundamental role in our construction of non-abelian local class field theory [9], which generalizes Laubie theory [11] as well.

Acknowledgements

The first named author (K. I. I.) would like to thank to the Institut de Mathématiques, “Théorie des Groupes, Représentations et Applications”, Université Paris 7, Jussieu, Paris, France, and to the School of Pure Mathematics of the Tata Institute of Fundamental Research, Mumbai, India, for their hospitality and support, where some parts of this work has been completed. The second named author (E. S.) would like to thank to TÜBITAK for a fellowship, and to the School of Mathematics of the University of Nottingham for the hospitality and support, where some parts of this work have been introduced. Both of the authors would like to thank I. B. Fesenko for his interest and encouragement at all stages of this work.

Notation


All through this work, K will denote a local field (a complete discrete valuation field) with finite residue field $O_K/\mathfrak{p}_K =: \kappa_K$ of $q_K = q = p^f$ elements with p a prime number, where O_K denotes the ring of integers in K with the unique maximal ideal $\mathfrak{p}_K$. Let ν_K denote the corresponding normalized valuation on K (normalized by $\nu_K(K^\times) = \mathbb{Z}$), and $\tilde{\nu}$ the unique extension of ν_K to a fixed separable closure K^{sep} of K . For any sub-extension L/K of K^{sep}/K , the normalized form of the valuation $\tilde{\nu}|_L$ on L will be denoted by ν_L . As usual, we let K^{nr} to denote the maximal unramified extension in K^{sep} , and $\tilde{K}$ denotes the completion of K^{nr} with respect to $\nu_{K^{nr}}$. Fix a *Lubin-Tate splitting* $\varphi_K = \varphi$ over K . The fixed field of the Lubin-Tate splitting φ is denoted by K_φ . Finally, let $(\pi_E)_{K \subset E \subset K_\varphi}$ be the canonical sequence of norm-compatible prime elements in finite sub-extensions E/K in K_φ/K . This determines a unique *Lubin-Tate labelling over K* (cf. paragraph 0.2 of [10]).

1 Preliminaries on Fontaine-Wintenberger field of norms

For a brief review of *APF*-extensions and Fontaine-Wintenberger field of norms, we refer the reader to [8], and for detailed proofs to [5, 6, 12].

Let L/K be an infinite Galois arithmetically profinite (in short *APF*) extension such that the residue-class degree $[\kappa_L : \kappa_K] = d$ and $K \subset L \subset K_{\varphi^d}$; that is, in the terminology of Koch-de Shalit in [10] and Laubie in [11], L is *compatible with (T, φ)* , where T denotes the intersection field $L \cap K_\varphi$. Note that, T/K is in general *not* a normal extension! Denote by $L_0^{(K)} = L \cap K^{nr} = K_d^{nr}$. If there is no fear of confusion, denote $L_0^{(K)}$ simply by L_0 . So, there is the following

diagram :



Remark 1.1. Note that, $\varphi' = \varphi^d$ is a Lubin-Tate splitting over $L_0 = K_d^{nr}$. Thus, by Proposition 1.2.3 of [12] or by Lemma 3.3 of [8], L/L_0 is an infinite totally-ramified APF-Galois extension satisfying $L_0 \subseteq L \subseteq (L_0)_{\varphi'}$. Thus, Fesenko theory, developed in [1, 2, 3] and [8], works for the extension L/L_0 .

Since L/T is an unramified extension, we have the following lemma.

Lemma 1.2. *The field of norms $\mathbb{X}(L/L_0)$ is an unramified extension of the field of norms $\mathbb{X}(T/K)$.*

Proof. In fact, there exists a natural isomorphism $\mathbb{X}(L/L_0) \xrightarrow{\sim} \mathbb{X}(L/K)$ which identifies $\mathbb{X}(L/L_0)$ and $\mathbb{X}(L/K)$ (cf. section (5.6) of Chapter III of [4]). Now, $\mathbb{X}(L/K)$ is a Galois extension of $\mathbb{X}(T/K)$ with corresponding Galois group isomorphic to $\text{Gal}(L/T)$ (cf. [8] and [12]). As $\kappa_{\mathbb{X}(L/K)} \simeq \kappa_L$ and $\kappa_{\mathbb{X}(T/K)} \simeq \kappa_T$, it follows that

$$[\kappa_{\mathbb{X}(L/K)} : \kappa_{\mathbb{X}(T/K)}] = [\mathbb{X}(L/K) : \mathbb{X}(T/K)],$$

as L/T is an unramified extension, which proves that $\mathbb{X}(L/K)$ is an unramified extension of $\mathbb{X}(T/K)$. $\square$

Now, as the Lubin-Tate splitting φ over K is fixed, the unique element $\Pi_{\varphi;T/K} = (\pi_E)_{K \subset E \subset T} \in \mathbb{X}(T/K)$ is a canonical prime element of the local field $\mathbb{X}(T/K)$. Thus, in view of Lemma 1.2, $\Pi_{\varphi;T/K}$ is a prime element of $\mathbb{X}(L/L_0)$ as well. Moreover,

Lemma 1.3.

$$\Pi_{\varphi;T/K} = \Pi_{\varphi';L/L_0}. \quad (1.1)$$

Proof. In fact, for the Lubin-Tate splitting $\varphi' = \varphi^d$ over $L_0 = K_d^{nr}$, there exists a unique element $(\pi_{L_0 E})_{L_0 \subset L_0 E \subset L_0 T = L} \in \mathbb{X}(L/L_0)$. As $L_0 E/E$ is an unramified extension, it follows that $\pi_{L_0 E} = \pi_E$ for each $K \subset E \subset T$. Thus, eq. (1.1) follows. $\square$

The completion $\widetilde{\mathbb{X}}(L/K)$ of the maximal unramified extension $\mathbb{X}(L/K)^{nr}$ of the field of norms $\mathbb{X}(L/K)$ is identified with the field of norms $\mathbb{X}(\widetilde{L}/\widetilde{K}) = \mathbb{X}(\widetilde{L}/\widetilde{L_0})$.

2 Generalized Fesenko reciprocity map

The main references for this section are [1, 2, 3] and [8].

Fix a Lubin-Tate splitting $\varphi_K = \varphi$ over K . The aim of this section is to generalize the reciprocity map $\Phi_{M/K}^{(\varphi)}$ of Fesenko, cf. [1, 2, 3] and [8], defined for infinite totally-ramified *APF*-Galois extensions M/K satisfying $K \subset M \subset K_\varphi$ to infinite *APF*-Galois extensions L/K with residue-class degree $[\kappa_L : \kappa_K] = d$ and satisfying $K \subset L \subset K_{\varphi^d}$. We shall keep the notation introduced in [8] and introduced in the previous section in what follows.

Recall that, for the extension M/K as above, the diamond subgroup $U_{\tilde{\mathbb{X}}(M/K)}^\diamond$ of the group $U_{\tilde{\mathbb{X}}(M/K)}$ of units of the ring of integers of $\tilde{\mathbb{X}}(M/K)$ is defined by

$$U_{\tilde{\mathbb{X}}(M/K)}^\diamond = \text{Pr}_{\tilde{K}}^{-1}(U_K),$$

where $\text{Pr}_{\tilde{K}} : U_{\tilde{\mathbb{X}}(M/K)} \rightarrow U_{\tilde{K}}$ denotes the projection map on the $\tilde{K}$ -coordinate of $U_{\tilde{\mathbb{X}}(M/K)}$. More generally, for a given infinite *APF*-Galois extension L/K with residue-class degree $[\kappa_L : \kappa_K] = d$ and satisfying $K \subset L \subset K_{\varphi^d}$, define the diamond subgroup $U_{\tilde{\mathbb{X}}(L/K)}^\diamond$ of the group $U_{\tilde{\mathbb{X}}(L/K)}$ of units of the ring of integers of $\tilde{\mathbb{X}}(L/K) = \tilde{\mathbb{X}}(L/L_0)$ naturally as follows.

Definition 2.1. $U_{\tilde{\mathbb{X}}(L/K)}^\diamond$ is the subgroup of the group $U_{\tilde{\mathbb{X}}(L/K)}$ of units of the ring of integers of the local field $\tilde{\mathbb{X}}(L/K)$ whose $\tilde{K} = \tilde{L}_0$ -coordinate belongs to U_{L_0} . That is,

$$U_{\tilde{\mathbb{X}}(L/K)}^\diamond = U_{\tilde{\mathbb{X}}(L/L_0)}^\diamond.$$

Now, as a first step, we shall generalize the arrow $\phi_{M/K}^{(\varphi)}$ defined for the extensions M/K , where M/K is a totally-ramified *APF*-Galois extension satisfying $K \subset M \subset K_\varphi$, of Fesenko theory, which has been described in [1, 2, 3] and in detail in Section 5 of [8], to infinite *APF*-Galois extensions L of K with residue-class degree d and satisfying $K_d^{nr} \subset L \subset K_{\varphi^d}$ and construct the *generalized arrow* $\phi_{L/K}^{(\varphi)}$ for such extensions L/K as follows. There exists an isomorphism

$$\text{Gal}(L/K) \xrightarrow{\sim} \text{Gal}(L_0/K) \times \text{Gal}(L/L_0) \quad (2.1)$$

defined by

$$\sigma \mapsto (\sigma|_{L_0}, \varphi^{-m}\sigma), \quad (2.2)$$

for every $\sigma \in \text{Gal}(L/K)$, where $\sigma|_{L_0} = \varphi^m$ for some $0 \leq m \in \mathbb{Z}$.

Remark 2.2. (i) Let M/K be a Galois sub-extension of L/K . Let $M_0 = M \cap K^{nr}$. Then, the following square

$$\begin{array}{ccc} \text{Gal}(L/K) & \xrightarrow{\sim} & \text{Gal}(L_0/K) \times \text{Gal}(L/L_0) \\ \text{res}_M \downarrow & & \downarrow (\text{res}_{M_0}, \text{res}_M) \\ \text{Gal}(M/K) & \xrightarrow{\sim} & \text{Gal}(M_0/K) \times \text{Gal}(M/M_0) \end{array}$$

is commutative. Now, for $\sigma \in \text{Gal}(L/K)$, there exists $0 \leq m, m' \in \mathbb{Z}$ such that $\sigma|_{L_0} = \varphi^m$ and $(\sigma|_M)|_{M_0} = \varphi^{m'}$. Thus, $\varphi^m|_{M_0} = \varphi^{m'}|_{M_0}$ and the identity $(\varphi^{-m}\sigma)|_M = \varphi^{-m'}(\sigma|_M)$ is satisfied.

- (ii) Let F/K be a finite Galois sub-extension of L/K . Let $L_0^{(K)} = L \cap K^{nr}$ and $L_0^{(F)} = L \cap F^{nr}$. Then, the following square

$$\begin{array}{ccc} \mathrm{Gal}(L/F) & \xrightarrow{\sim} & \mathrm{Gal}(L_0^{(F)}/F) \times \mathrm{Gal}(L/L_0^{(F)}) \\ \mathrm{inc.} \downarrow & & \downarrow (res_{L_0^{(K)}}, \mathrm{inc.}) \\ \mathrm{Gal}(L/K) & \xrightarrow{\sim} & \mathrm{Gal}(L_0^{(K)}/K) \times \mathrm{Gal}(L/L_0^{(K)}) \end{array}$$

is commutative. Now, for any $\sigma \in \mathrm{Gal}(L/F)$, there exists $0 \leq m, m' \in \mathbb{Z}$ such that $\sigma|_{L_0^{(F)}} = \varphi_F^m$ and $\sigma|_{L_0^{(K)}} = \varphi_K^{m'}$. Thus, $\varphi_F^m|_{L_0^{(K)}} = \varphi_K^{m'}$ and the identity $\varphi_F^{-m}\sigma = \varphi_K^{-m'}\sigma$ is satisfied.

Now, by Proposition 1.2.3 of [12] or by Lemma 3.3 of [8], L/L_0 is a totally-ramified *APF*-Galois extension with $L_0 \subset L \subset (L_0)_{\varphi'}$, where $\varphi' = \varphi^d$ is a Lubin-Tate splitting over L_0 by Remark 1.1. Thus, define the map

$$\phi_{L/K}^{(\varphi)} : \mathrm{Gal}(L/K) \rightarrow K^\times / N_{L_0/K} L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)} \quad (2.3)$$

by

$$\phi_{L/K}^{(\varphi)}(\sigma) := (\pi_K^m \cdot N_{L_0/K} L_0^\times, (u_{\tilde{E}}) \cdot U_{\mathbb{X}(L/K)}), \quad (2.4)$$

where $\sigma \in \mathrm{Gal}(L/K)$ such that $\sigma|_{L_0} = \varphi^m$, for some $0 \leq m \in \mathbb{Z}$, and $U = (u_{\tilde{E}}) \in U_{\mathbb{X}(L/L_0)}^\diamond$ satisfies the equality

$$U^{1-\varphi^d} = \Pi_{\varphi'; L/L_0}^{(\varphi^{-m}\sigma)^{-1}}, \quad (2.5)$$

where $\Pi_{\varphi'; L/L_0}$ is the canonical prime element of the local field $\mathbb{X}(L/L_0)$, which is the canonical prime element $\Pi_{\varphi; T/K}$ of the local field $\mathbb{X}(T/K)$ by Lemma 1.2 and by Lemma 1.3. Thus, (2.5) can be reformulated by

$$U^{1-\varphi^d} = \Pi_{\varphi'; L/L_0}^{\sigma^{-1}}, \quad (2.6)$$

as $\Pi_{\varphi'; T/K} = \Pi_{\varphi; T/K}$. Moreover, the solution $U = (u_{\tilde{E}}) \in U_{\mathbb{X}(L/L_0)}^\diamond$, which is unique modulo $U_{\mathbb{X}(L/K)}$, satisfies $\mathrm{Pr}_{L_0}(U) = u_{\tilde{L}_0} \in U_{L_0}$. In fact, by Lemma 1.3, $\mathrm{Pr}_{L_0}(\Pi_{\varphi'; L/L_0}) = \pi_K$, and therefore $\mathrm{Pr}_{L_0}(\Pi_{\varphi'; L/L_0}^{\sigma^{-1}}) = \pi_K^{\sigma^{-1}} = 1_K$. Hence, $\mathrm{Pr}_{L_0}(U^{1-\varphi^d}) = \mathrm{Pr}_{L_0}(\Pi_{\varphi'; L/L_0}^{\sigma^{-1}}) = 1_K$ yields $u_{\tilde{L}_0}^{1-\varphi^d} = 1_K$, that is $u_{\tilde{L}_0} \in U_{L_0}$ as $\tilde{L}_0 \cap (L_0)_{\varphi'} = L_0$. Now, it follows that, $\mathrm{Pr}_{L_0}(U) = u_{\tilde{L}_0} \in U_{L_0}$. Thus, $U = (u_{\tilde{E}})$ belongs to $U_{\mathbb{X}(L/K)}^\diamond$, by Definition 2.1.

Remark 2.3. We can reformulate the definition of the generalized arrow

$$\phi_{L/K}^{(\varphi)} : \mathrm{Gal}(L/K) \rightarrow K^\times / N_{L_0/K} L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)}$$

for the extension L/K as

$$\phi_{L/K}^{(\varphi)}(\sigma) = \left(\pi_K^m \cdot N_{L_0/K} L_0^\times, \phi_{L/L_0}^{(\varphi')}(\varphi^{-m}\sigma) \right)$$

for every $\sigma \in \mathrm{Gal}(L/K)$, where $\sigma|_{L_0} = \varphi^m$ for some $0 \leq m \in \mathbb{Z}$.

There is a natural continuous action of $\text{Gal}(L/K)$ on the topological group $K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)}$ defined by abelian local class field theory on the first component and by eq.s (5.5) and (5.7) of [8] on the second component as

$$(\bar{a}, \bar{U})^\sigma = \left(\bar{a}^{\varphi^m}, \bar{U}^{\varphi^{-m}\sigma} \right) = \left(\bar{a}, \bar{U}^{\varphi^{-m}\sigma} \right), \quad (2.7)$$

for every $\sigma \in \text{Gal}(L/K)$, where $\sigma|_{L_0} = \varphi^m$ for some $0 \leq m \in \mathbb{Z}$, and for every $a \in K^\times$ with $\bar{a} = a.N_{L_0/K}L_0^\times$ and $U \in U_{\mathbb{X}(L/K)}^\diamond$ with $\bar{U} = U.U_{\mathbb{X}(L/K)}$. We shall always view $K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)}$ as a topological $\text{Gal}(L/K)$ -module in this text.

Theorem 2.4. *Let L/K be any infinite APF-Galois sub-extension of K_{φ^d}/K with residue-class degree d . Then the generalized arrow*

$$\phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \rightarrow K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)}$$

defined for the extension L/K is an injection, and for every $\sigma, \tau \in \text{Gal}(L/K)$,

$$\phi_{L/K}^{(\varphi)}(\sigma\tau) = \phi_{L/K}^{(\varphi)}(\sigma)\phi_{L/K}^{(\varphi)}(\tau)^\sigma \quad (2.8)$$

co-cycle condition is satisfied.

Proof. The injectivity of the arrow given by eq. (2.3) and defined by eq. (2.4) is clear from the canonical topological isomorphism defined by (2.1) and (2.2) combined with abelian local class field theory and Theorem 5.6 of Fesenko in [8]. To be precise, let $\phi_{L/K}^{(\varphi)}(\sigma) = (\pi_K^m, (u_{\bar{E}}))$ with $d \mid m$ and $(u_{\bar{E}}) \in U_{\mathbb{X}(L/L_0)} = U_{\mathbb{X}(L/K)}$. As $d \mid m$, σ acts trivially on L_0 . Since $(u_{\bar{E}})^{\varphi^d-1} = (1_{\bar{E}}) = \Pi_{\varphi'; L/L_0}^{\sigma-1}$, σ acts trivially on the prime elements of finite sub-extensions between L_0 and L . Thus, σ is the identity element of $\text{Gal}(L/L_0)$. Now, for $\sigma, \tau \in \text{Gal}(L/K)$, with $\sigma|_{L_0} = \varphi^m$ and $\tau|_{L_0} = \varphi^n$ for some $0 \leq m, n \in \mathbb{Z}$, following the alternative definition of the generalized arrow $\phi_{L/K}^{(\varphi)}$ introduced in Remark 2.3,

$$\begin{aligned} \phi_{L/K}^{(\varphi)}(\sigma\tau) &= (\pi_K^{m+n}.N_{L_0/K}L_0^\times, \phi_{L/L_0}^{(\varphi')}(\varphi^{-(m+n)}\sigma\tau)) \\ &= \left((\pi_K^m.N_{L_0/K}L_0^\times)(\pi_K^n.N_{L_0/K}L_0^\times), \phi_{L/L_0}^{(\varphi')}(\varphi^{-m}\sigma)\phi_{L/L_0}^{(\varphi')}(\varphi^{-n}\tau)^{\varphi^{-m}\sigma} \right) \\ &= \left(\pi_K^m.N_{L_0/K}L_0^\times, \phi_{L/L_0}^{(\varphi')}(\varphi^{-m}\sigma) \right) \left(\pi_K^n.N_{L_0/K}L_0^\times, \phi_{L/L_0}^{(\varphi')}(\varphi^{-n}\tau)^{\varphi^{-m}\sigma} \right) \\ &= \phi_{L/K}^{(\varphi)}(\sigma)\phi_{L/K}^{(\varphi)}(\tau)^\sigma \end{aligned}$$

by Theorem 5.6 of Fesenko in [8] and by the definition of the action of $\sigma \in \text{Gal}(L/K)$ on $\phi_{L/K}^{(\varphi)}(\tau) \in K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)}$ defined by eq. (2.7). $\square$

Now, we immediately have the following result.

Corollary 2.5. *Define a law of composition $*$ on $\text{im}(\phi_{L/K}^{(\varphi)})$ by*

$$(\bar{a}, \bar{U}) * (\bar{b}, \bar{V}) = (\bar{a}.\bar{b}, \bar{U}.\bar{V}^{(\phi_{L/L_0}^{(\varphi')})^{-1}(\bar{U})}), \quad (2.9)$$

for $(\bar{a}, \bar{U}), (\bar{b}, \bar{V}) \in \text{im}(\phi_{L/K}^{(\varphi)})$, where $\bar{a} = a.N_{L_0/K}L_0^\times, \bar{b} = b.N_{L_0/K}L_0^\times \in K^\times/N_{L_0/K}L_0^\times$ with $a, b \in K^\times$ and for $\bar{U} = U.U_{\mathbb{X}(L/K)}, \bar{V} = V.U_{\mathbb{X}(L/K)} \in U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)}$ with $U, V \in U_{\mathbb{X}(L/K)}^\diamond$. Then $\text{im}(\phi_{L/K}^{(\varphi)})$ is a topological group under $*$, and the map $\phi_{L/K}^{(\varphi)}$ induces an isomorphism of topological groups

$$\phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \xrightarrow{\sim} \text{im}(\phi_{L/K}^{(\varphi)}), \quad (2.10)$$

where the topological group structure on $\text{im}(\phi_{L/K}^{(\varphi)})$ is defined with respect to the binary operation $*$ defined by eq. (2.9).

Recall that, for an infinite APF-Galois extension L/K and for every $-1 \leq u \in \mathbb{R}$, the u^{th} higher ramification subgroup $\text{Gal}(L/K)_u$ of $\text{Gal}(L/K)$ in lower numbering is defined by

$$\text{Gal}(L/K)_u = \text{Gal}(L/K)^{\varphi_{L/K}(u)},$$

where the number $-1 \leq \varphi_{L/K}(u) \in \mathbb{R}$ is defined by eq. (3.1) in [8] and as usual the $\varphi_{L/K}(u)^{\text{th}}$ higher ramification subgroup $\text{Gal}(L/K)^{\varphi_{L/K}(u)}$ of $\text{Gal}(L/K)$ in upper numbering by the projective limit $\text{Gal}(L/K)^{\varphi_{L/K}(u)} = \varprojlim_{K \subseteq F \subset L} \text{Gal}(F/K)^{\varphi_{L/K}(u)}$ defined by eq.s (2.1) and (2.2) in [8]. Now, let E/K be a Galois sub-extension

of L/K . Then, for any chain of field extensions $\underbrace{K \subseteq F \subseteq F'}_{\text{finite Gal.}} \subset L$, the square

$$\begin{array}{ccc} \text{Gal}(F'/K)^{\varphi_{L/K}(u)} & \xrightarrow{t_{F' \cap E}^{F'}(\varphi_{L/K}(u))} & \text{Gal}(F' \cap E/K)^{\varphi_{L/K}(u)} \\ \downarrow t_F^{F'}(\varphi_{L/K}(u)) & & \downarrow t_{F \cap E}^{F' \cap E}(\varphi_{L/K}(u)) \\ \text{Gal}(F/K)^{\varphi_{L/K}(u)} & \xrightarrow{t_{F \cap E}^F(\varphi_{L/K}(u))} & \text{Gal}(F \cap E/K)^{\varphi_{L/K}(u)} \end{array} \quad (2.11)$$

is commutative. Thus, passing to the projective limits, there exists a continuous group homomorphism

$$t_E^L(\varphi_{L/K}(u)) = \varprojlim_{K \subseteq F \subset L} t_{F \cap E}^F(\varphi_{L/K}(u)) : \text{Gal}(L/K)^{\varphi_{L/K}(u)} \rightarrow \text{Gal}(E/K)^{\varphi_{L/K}(u)}, \quad (2.12)$$

which is essentially the restriction morphism from L to E . This morphism is a surjection, as the objects in the respective projective systems are compact and Hausdorff. Furthermore, the following square

$$\begin{array}{ccc} \text{Gal}(L/K) & \xrightarrow{r_E^L} & \text{Gal}(E/K) \\ \uparrow \text{inc.} & & \uparrow \text{inc.} \\ \text{Gal}(L/K)^{\varphi_{L/K}(u)} & \xrightarrow{t_E^L(\varphi_{L/K}(u))} & \text{Gal}(E/K)^{\varphi_{L/K}(u)} \\ \uparrow \text{inc.} & & \uparrow \text{inc.} \\ \text{Gal}(L/K)^{\varphi_{L/K}(u')} & \xrightarrow{t_E^L(\varphi_{L/K}(u'))} & \text{Gal}(E/K)^{\varphi_{L/K}(u')} \end{array} \quad (2.13)$$

is commutative for every pair $u, u' \in \mathbb{R}_{\geq -1}$ satisfying $u \leq u'$. Here, the arrow $r_E^L : \text{Gal}(L/K) \rightarrow \text{Gal}(E/K)$ denotes the restriction map. Therefore, immediately, we have the following result.

Lemma 2.6. *For $0 \leq u \in \mathbb{R}$, the topological isomorphism defined by eq.s (2.1) and (2.2) induces a topological isomorphism*

$$\text{Gal}(L/K)_u \simeq \underbrace{\text{Gal}(L_0/K)^{\varphi_{L/K}(u)}}_{\langle \text{id}_{L_0} \rangle} \times \text{Gal}(L/L_0)^{\varphi_{L/K}(u)} \quad (2.14)$$

defined by

$$\sigma \mapsto (t_{L_0}^L(\varphi_{L/K}(u))(\sigma), \varphi^{-m}\sigma) = (\text{id}_{L_0}, \sigma), \quad (2.15)$$

for every $\sigma \in \text{Gal}(L/K)_u$ with $\sigma|_{L_0} = t_{L_0}^L(\varphi_{L/K}(u))(\sigma) = \varphi^m|_{L_0}$ for some $0 \leq m \in \mathbb{Z}$ satisfying $d \mid m$.

Proof. Note that, $\text{Gal}(L_0/K)^{\varphi_{L/K}(u)}$, for $0 \leq u \in \mathbb{R}$, is the trivial group $< \text{id}_{L_0} >$, as L_0/K is a finite unramified extension. Thus, for $\sigma \in \text{Gal}(L/K)_u$, by the commutativity of the diagram (2.13),

$$t_{L_0}^L(\varphi_{L/K}(u))(\sigma) = \sigma|_{L_0} = \text{id}_{L_0},$$

and in return

$$\sigma \mapsto (\text{id}_{L_0}, \varphi^{-m}\sigma) = (\text{id}_{L_0}, \sigma),$$

where $\sigma|_{L_0} = \text{id}_{L_0} = \varphi^m|_{L_0}$ for some $0 \leq m \in \mathbb{Z}$ satisfying $d \mid m$. As L is fixed by φ^d , $\varphi^{-m}\sigma = \sigma$, thus $(\text{id}_{L_0}, \varphi^{-m}\sigma) = (\text{id}_{L_0}, \sigma)$. Now, the injectivity of the morphism given by eq. (2.14) and defined by eq. (2.15) is clear from the commutative square eq. (2.13) and by the injectivity of the arrow given by eq. (2.1) and defined by eq. (2.2). Thus it suffices to prove that this morphism is a surjection, which follows from the triviality of $\text{Gal}(L_0/K)^{\varphi_{L/K}(u)}$ for $0 \leq u \in \mathbb{R}$, and from the equality $\text{Gal}(L/K)_u = \text{Gal}(L/L_0)^{\varphi_{L/K}(u)}$. $\square$

Now, L/L_0 is an *APF*-Galois sub-extension of L/K by part (i) of Lemma 3.3 in [8]. Let $\varphi_{L/L_0} : \mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$ be the Hasse-Herbrand function corresponding to the *APF*-extension L/L_0 defined by eq. (3.1) in [8], which is piecewise-linear and continuous. So there exists a unique number $w = w(u, L/K) \in \mathbb{R}_{\geq -1}$, depending on u , satisfying $\varphi_{L/K}(u) = \varphi_{L/L_0}(w)$ and

$$\text{Gal}(L/L_0)^{\varphi_{L/K}(u)} = \text{Gal}(L/L_0)^{\varphi_{L/L_0}(w)} = \text{Gal}(L/L_0)_w.$$

Thus, Lemma 2.6 can be reformulated as follows. The topological isomorphism defined by eq.s (2.1) and (2.2) induces a topological isomorphism

$$\text{Gal}(L/K)_u \simeq \langle \text{id}_{L_0} \rangle \times \text{Gal}(L/L_0)_{w(u, L/K)},$$

for every $0 \leq u \in \mathbb{R}$.

For each $0 \leq i \in \mathbb{R}$, consider the i^{th} higher unit group $U_{\tilde{\mathbb{X}}(L/K)}^i$ of the field $\tilde{\mathbb{X}}(L/K)$, and define the group

$$\left(U_{\tilde{\mathbb{X}}(L/K)}^\diamond \right)^i = U_{\tilde{\mathbb{X}}(L/K)}^\diamond \cap U_{\tilde{\mathbb{X}}(L/K)}^i. \quad (2.16)$$

Now, Fesenko ramification theorem, stated as Theorem 5.8 in [8], has the following generalization for the generalized arrow $\phi_{L/K}^{(\varphi)}$ corresponding to the extension L/K , which is an infinite APF-Galois sub-extension of K_{φ^d}/K with residue-class degree $[\kappa_L : \kappa_K] = d$.

Theorem 2.7 (Ramification theorem). *For $0 \leq u \in \mathbb{R}$, let $\text{Gal}(L/K)_u$ denote the u^{th} higher ramification subgroup in the lower numbering of the Galois group $\text{Gal}(L/K)$ corresponding to the infinite APF-Galois sub-extension L/K of K_{φ^d}/K with residue-class degree $[\kappa_L : \kappa_K] = d$. Then, for $0 \leq n \in \mathbb{Z}$, there exists the inclusion*

$$\phi_{L/K}^{(\varphi)} \left(\text{Gal}(L/K)_{\psi_{L/K} \circ \varphi_{L/L_0}(n)} - \text{Gal}(L/K)_{\psi_{L/K} \circ \varphi_{L/L_0}(n+1)} \right) \subseteq \left\langle 1_{K^\times/N_{L_0/K}L_0^\times} \right\rangle \times \left(\left(U_{\mathbb{X}(L/K)}^\circ \right)^n U_{\mathbb{X}(L/K)}/U_{\mathbb{X}(L/K)} - \left(U_{\mathbb{X}(L/K)}^\circ \right)^{n+1} U_{\mathbb{X}(L/K)}/U_{\mathbb{X}(L/K)} \right).$$

Proof. Now, we start with the following general observation. Let $0 \leq u \in \mathbb{R}$. Let $\tau \in \text{Gal}(L/K)_u = \text{Gal}(L/L_0)^{\varphi_{L/K}(u)}$. Then, by the definition of the generalized arrow $\phi_{L/K}^{(\varphi)}$ reformulated as in Remark 2.3,

$$\phi_{L/K}^{(\varphi)}(\tau) = \left(\pi_K^m \cdot N_{L_0/K} L_0^\times, \phi_{L/L_0}^{(\varphi')}(\varphi^{-m}\tau) \right),$$

where $\tau|_{L_0} = \varphi^m|_{L_0}$, for some $0 \leq m \in \mathbb{Z}$ satisfying $d \mid m$, as $\tau \in \text{Gal}(L/K)_u$ and $\tau|_{L_0} = t_{L_0}^L(\varphi_{L/K}(u))(\tau) \in \text{Gal}(L_0/K)^{\varphi_{L/K}(u)} = \langle \text{id}_{L_0} \rangle$. Thus,

$$\phi_{L/K}^{(\varphi)}(\tau) = \left(1_{K^\times/N_{L_0/K}L_0^\times}, \phi_{L/L_0}^{(\varphi')}(\varphi^{-m}\tau) \right),$$

as $m = dm'$ and thereby $\pi_K^{dm'} N_{L_0/K} L_0^\times = N_{L_0/K} L_0^\times = 1_{K^\times/N_{L_0/K}L_0^\times}$, since $N_{L_0/K} \pi_K^{m'} = \pi_K^m$. Therefore,

$$\phi_{L/K}^{(\varphi)}(\tau) = \left(1_{K^\times/N_{L_0/K}L_0^\times}, \phi_{L/L_0}^{(\varphi')}(\tau) \right),$$

since $\varphi^{-m}\tau = \tau$ in $\text{Gal}(L/L_0)$, as $d \mid m$, and $L \subset K_{\varphi^d}$.

Now, to prove the theorem, let $u = \psi_{L/K} \circ \varphi_{L/L_0}(n)$ and $u' = \psi_{L/K} \circ \varphi_{L/L_0}(n+1)$. Then, for any $\tau \in \text{Gal}(L/K)_u - \text{Gal}(L/K)_{u'}$, it follows from Fesenko ramification theorem (cf. Theorem 5.8 in [8]) that the second coordinate of $\phi_{L/K}^{(\varphi)}(\tau)$ satisfies

$$\phi_{L/L_0}^{(\varphi')}(\tau) \in \left(U_{\mathbb{X}(L/K)}^\circ \right)^n U_{\mathbb{X}(L/K)}/U_{\mathbb{X}(L/K)} - \left(U_{\mathbb{X}(L/K)}^\circ \right)^{n+1} U_{\mathbb{X}(L/K)}/U_{\mathbb{X}(L/K)},$$

since

$$\text{Gal}(L/K)_u = \text{Gal}(L/L_0)^{\varphi_{L/K}(u)} = \text{Gal}(L/L_0)^{\varphi_{L/L_0}(n)} = \text{Gal}(L/L_0)_n$$

and likewise

$$\text{Gal}(L/K)_{u'} = \text{Gal}(L/L_0)^{\varphi_{L/K}(u')} = \text{Gal}(L/L_0)^{\varphi_{L/L_0}(n+1)} = \text{Gal}(L/L_0)_{n+1},$$

which completes the proof. $\square$

Now, let M/K be an infinite Galois sub-extension of L/K . Thus, by Lemma 3.3 of [8], M is an *APF*-Galois extension over K . We further assume that, the residue-class degree $[\kappa_M : \kappa_K] = d'$ and $K \subset M \subset K_{\varphi^{d'}}$ for some $d' \mid d$. Let

$$\phi_{M/K}^{(\varphi)} : \text{Gal}(M/K) \rightarrow K^\times / N_{M_0/K} M_0^\times \times U_{\tilde{\mathbb{X}}(M/K)}^\diamond / U_{\mathbb{X}(M/K)}$$

be the corresponding generalized arrow defined for the extension M/K . Here, M_0 is defined by $M_0 = M \cap K^{nr} = K_{d'}^{nr}$.

Now, let

$$K \subset L_o = E_o \subset E_1 \subset \cdots \subset E_i \subset \cdots \subset L$$

be an ascending chain satisfying $L = \bigcup_{0 \leq i \in \mathbb{Z}} E_i$ and $[E_{i+1} : E_i] < \infty$ for every $0 \leq i \in \mathbb{Z}$. Then

$$K \subset M_o = E_o \cap M \subseteq E_1 \cap M \subseteq \cdots \subseteq E_i \cap M \subseteq \cdots \subset M$$

is an ascending chain of field extensions satisfying the conditions $M = \bigcup_{0 \leq i \in \mathbb{Z}} (E_i \cap M)$ and also $[E_{i+1} \cap M : E_i \cap M] < \infty$ for every $0 \leq i \in \mathbb{Z}$. Thus, we construct $\mathbb{X}(M/K)$ by the sequence $(E_i \cap M)_{0 \leq i \in \mathbb{Z}}$ and $\tilde{\mathbb{X}}(M/K)$ by the sequence $(\widetilde{E_i \cap M})_{0 \leq i \in \mathbb{Z}}$. Note that, $E_i \cap M \neq E_i$ for every $0 \leq i \in \mathbb{Z}$. Furthermore, the commutative square

$$\begin{array}{ccc} \widetilde{E_i}^\times & \xleftarrow{\tilde{N}_{E_{i'}/E_i}} & \widetilde{E_{i'}}^\times \\ \downarrow \Pi_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell \tilde{N}_{E_i/E_i \cap M} & & \downarrow \Pi_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell \tilde{N}_{E_{i'}/E_{i'} \cap M} \\ \widetilde{E_i \cap M}^\times & \xleftarrow{\tilde{N}_{E_{i'} \cap M/E_i \cap M}} & \widetilde{E_{i'} \cap M}^\times \end{array}$$

for every pair $0 \leq i, i' \in \mathbb{Z}$ satisfying $i \leq i'$, induces the group homomorphism

$$\tilde{\mathcal{N}}_{L/M} = \varprojlim_{0 \leq i \in \mathbb{Z}} \left(\prod_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell \tilde{N}_{E_i/E_i \cap M} \right) : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times \quad (2.17)$$

defined by

$$\tilde{\mathcal{N}}_{L/M} \left((\alpha_{\tilde{E_i}})_{0 \leq i \in \mathbb{Z}} \right) = \left(\prod_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell \tilde{N}_{E_i/E_i \cap M} (\alpha_{\tilde{E_i}}) \right)_{0 \leq i \in \mathbb{Z}}, \quad (2.18)$$

for every $(\alpha_{\tilde{E_i}})_{0 \leq i \in \mathbb{Z}} \in \tilde{\mathbb{X}}(L/K)^\times$.

Remark 2.8. The group homomorphism

$$\tilde{\mathcal{N}}_{L/M} : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times$$

defined by eq.s (2.17) and (2.18) does *not* depend on the choice of the ascending chain

$$K \subset L_o = E_o \subset E_1 \subset \cdots \subset E_i \subset \cdots \subset L$$

satisfying $L = \bigcup_{0 \leq i \in \mathbb{Z}} E_i$ and $[E_{i+1} : E_i] < \infty$ for every $0 \leq i \in \mathbb{Z}$.

Remark 2.9. For $0 \leq i \in \mathbb{Z}$, let $E_{i,0}^{(E_i \cap M)} = E_i \cap (E_i \cap M)^{nr}$ be the maximal unramified extension of $E_i \cap M$ inside E_i . To simplify the notation, let $E_{i,0} = E_i \cap (E_i \cap M)^{nr}$. Then the Galois group $\text{Gal}(E_{i,0}/E_i \cap M)$ is cyclic of order $f(L/M) = \frac{d}{d'}$ generated by $\varphi^{d'}$. Thus, for $\alpha \in E_i$,

$$N_{E_i/E_i \cap M}(\alpha) = \tilde{N}_{E_i/E_i \cap M}(\alpha)^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}}.$$

The basic properties of this group homomorphism are the following.

(i) If $U = (u_{\tilde{E}_i})_{0 \leq i \in \mathbb{Z}} \in U_{\tilde{\mathbb{X}}(L/K)}$, then $\tilde{\mathcal{N}}_{L/M}(U) \in U_{\tilde{\mathbb{X}}(M/K)}$.

Proof. In fact, following the definition of the valuation $\nu_{\tilde{\mathbb{X}}(M/K)}$ of $\tilde{\mathbb{X}}(M/K)$ and the definition of the valuation $\nu_{\tilde{\mathbb{X}}(L/K)}$ of $\tilde{\mathbb{X}}(L/K)$, it follows that

$$\begin{aligned} \nu_{\tilde{\mathbb{X}}(M/K)}\left(\tilde{\mathcal{N}}_{L/M}(U)\right) &= \nu_{\tilde{\mathbb{X}}(M/K)}\left(\left(\prod_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell \tilde{N}_{E_i/E_i \cap M}(u_{\tilde{E}_i})\right)_{0 \leq i \in \mathbb{Z}}\right) \\ &= \nu_{\tilde{K}}\left(\prod_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell (u_{\tilde{K}})\right) \\ &= \sum_{0 \leq \ell \leq f(L/M)} \nu_{\tilde{K}}\left((\varphi^{d'})^\ell (u_{\tilde{K}})\right) \\ &= \sum_{0 \leq \ell \leq f(L/M)} \nu_{\tilde{K}}(u_{\tilde{K}}) \\ &= 0, \end{aligned}$$

as $\nu_{\tilde{K}}\left((\varphi^{d'})^\ell (u_{\tilde{K}})\right) = \nu_{\tilde{K}}(u_{\tilde{K}})$ for $\ell = 1, \dots, f(L/M) - 1$, and

$$\nu_{\tilde{\mathbb{X}}(L/K)}(U) = \nu_{\tilde{K}}(u_{\tilde{K}}) = 0,$$

since $U \in U_{\tilde{\mathbb{X}}(L/K)}$. □

(ii) If $U = (u_{\tilde{E}_i})_{0 \leq i \in \mathbb{Z}} \in U_{\tilde{\mathbb{X}}(L/K)}^\diamond$, then $\tilde{\mathcal{N}}_{L/M}(U) \in U_{\tilde{\mathbb{X}}(M/K)}^\diamond$.

Proof. Note that, $\tilde{L}_0 = \tilde{K}$ and $\tilde{M}_0 = \tilde{K}$. Now, the assertion follows by observing that

$$\text{Pr}_{\tilde{K}}(U) = u_{\tilde{K}} \in U_{L_0}$$

and

$$\begin{aligned} \text{Pr}_{\tilde{K}}\left(\tilde{\mathcal{N}}_{L/M}(U)\right) &= \prod_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell \tilde{N}_{E_o/E_o \cap M}(u_{\tilde{E}_o}) \\ &= \prod_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell u_{\tilde{K}} \\ &= N_{E_o/E_o \cap M}(u_{\tilde{K}}) \in U_{M_0}. \end{aligned}$$

□

(iii) If $U = (u_{E_i})_{0 \leq i \in \mathbb{Z}} \in U_{\mathbb{X}(L/K)}$, then $\tilde{\mathcal{N}}_{L/M}(U) \in U_{\mathbb{X}(M/K)}$.

Proof. The assertion follows by the definition eq. (2.18) of the homomorphism eq. (2.17) combined with the fact that $\tilde{N}_{E_i/E_i \cap M}(u_{E_i})^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}} = N_{E_i/E_i \cap M}(u_{E_i})$ for every $u_{E_i} \in U_{E_i}$ and for every $0 \leq i \in \mathbb{Z}$ by Remark 2.9. $\square$

Note that, $\tilde{N}_{E_i/E_{i-1}}(\alpha^{1+\varphi^{d'}+\dots+\varphi^{d'(f-1)}}) = \tilde{N}_{E_i/E_{i-1}}(\alpha)^{1+\varphi^{d'}+\dots+\varphi^{d'(f-1)}}$, for any $\alpha \in \tilde{E}_i$ with $1 \leq i \in \mathbb{Z}$, where $f = f(L/M)$. Thus, there exists a homomorphism

$$\langle \varphi \rangle_{L/M} : \tilde{\mathbb{X}}(L/L_0)^\times \rightarrow \tilde{\mathbb{X}}(L/L_0)^\times \quad (2.19)$$

defined by

$$\langle \varphi \rangle_{L/M} : \left(\alpha_{\tilde{E}_i} \right)_{0 \leq i \in \mathbb{Z}} \mapsto \left(\alpha_{\tilde{E}_i}^{1+\varphi^{d'}+\dots+\varphi^{d'(f-1)}} \right)_{0 \leq i \in \mathbb{Z}}, \quad (2.20)$$

for every $(\alpha_{\tilde{E}_i})_{0 \leq i \in \mathbb{Z}} \in \tilde{\mathbb{X}}(L/L_0)^\times$. The basic properties of this group homomorphism are the following.

$$(i) \quad \langle \varphi \rangle_{L/M} \left(U_{\tilde{\mathbb{X}}(L/L_0)} \right) \subseteq U_{\tilde{\mathbb{X}}(L/L_0)}.$$

Proof. In fact, following the definition of the valuation $\nu_{\tilde{\mathbb{X}}(L/L_0)}$ of $\tilde{\mathbb{X}}(L/L_0)$, it follows that

$$\begin{aligned} \nu_{\tilde{\mathbb{X}}(L/L_0)} \left(\langle \varphi \rangle_{L/M}(U) \right) &= \nu_{\tilde{\mathbb{X}}(L/L_0)} \left(\left(\alpha_{\tilde{E}_i}^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}} \right)_{0 \leq i \in \mathbb{Z}} \right) \\ &= \nu_{\tilde{K}} \left(u_{\tilde{K}}^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}} \right) \\ &= \sum_{0 \leq \ell \leq f(L/M)} \nu_{\tilde{K}} \left((\varphi^{d'})^\ell(u_{\tilde{K}}) \right) \\ &= \sum_{0 \leq \ell \leq f(L/M)} \nu_{\tilde{K}}(u_{\tilde{K}}) \\ &= 0, \end{aligned}$$

as $\nu_{\tilde{K}}((\varphi^{d'})^\ell(u_{\tilde{K}})) = \nu_{\tilde{K}}(u_{\tilde{K}})$ for $\ell = 1, \dots, f(L/M) - 1$, and

$$\nu_{\tilde{\mathbb{X}}(L/K)}(U) = \nu_{\tilde{K}}(u_{\tilde{K}}) = 0,$$

since $U \in U_{\tilde{\mathbb{X}}(L/K)}$. $\square$

$$(ii) \quad \langle \varphi \rangle_{L/M} \left(U_{\tilde{\mathbb{X}}(L/L_0)}^\diamond \right) \subseteq U_{\tilde{\mathbb{X}}(L/L_0)}^\diamond.$$

Proof. Note that, $\tilde{L}_0 = \tilde{K}$. Now, the assertion follows by observing that

$$\text{Pr}_{\tilde{K}}(U) = u_{\tilde{K}} \in U_{L_0}$$

and

$$\begin{aligned}\Pr_{\tilde{K}}\left(\langle\varphi\rangle_{L/M}(U)\right) &= u_{\tilde{K}}^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}} \\ &= \prod_{0 \leq \ell \leq f(L/M)} (\varphi^{d'})^\ell u_{\tilde{K}} \\ &= N_{E_o/E_o \cap M}(u_{\tilde{K}}) \in U_{M_0} \subseteq U_{L_0}.\end{aligned}$$

□

$$(iii) \quad \langle\varphi\rangle_{L/M}(U_{\mathbb{X}(L/L_0)}) \subseteq U_{\mathbb{X}(L/L_0)}.$$

Proof. Clearly, for $U = (u_{E_i})_{0 \leq i \in \mathbb{Z}} \in U_{\mathbb{X}(L/L_0)}$,

$$\langle\varphi\rangle_{L/M}(U) = (u_{E_i}^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}})_{0 \leq i \in \mathbb{Z}} \in U_{\mathbb{X}(L/L_0)},$$

as $u_{E_i} \in U_{E_i}$ for every $0 \leq i \in \mathbb{Z}$. □

Thus, there exists a group homomorphism

$$\tilde{\mathcal{N}}_{L/M} \circ \langle\varphi\rangle_{L/M} : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times \quad (2.21)$$

satisfying

$$(i) \quad \tilde{\mathcal{N}}_{L/M} \circ \langle\varphi\rangle_{L/M} \left(U_{\tilde{\mathbb{X}}(L/K)} \right) \subseteq U_{\tilde{\mathbb{X}}(M/K)};$$

$$(ii) \quad \tilde{\mathcal{N}}_{L/M} \circ \langle\varphi\rangle_{L/M} \left(U_{\tilde{\mathbb{X}}(L/K)}^\diamond \right) \subseteq U_{\tilde{\mathbb{X}}(M/K)}^\diamond;$$

$$(iii) \quad \tilde{\mathcal{N}}_{L/M} \circ \langle\varphi\rangle_{L/M} \left(U_{\mathbb{X}(L/K)} \right) \subseteq U_{\mathbb{X}(M/K)}.$$

Now, define the *Coleman norm map*

$$\tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} : U_{\tilde{\mathbb{X}}(L/K)}^\diamond / U_{\mathbb{X}(L/K)} \rightarrow U_{\tilde{\mathbb{X}}(M/K)}^\diamond / U_{\mathbb{X}(M/K)} \quad (2.22)$$

from L to M by

$$\tilde{\mathcal{N}}_{L/M}^{\text{Coleman}}(\overline{U}) = \tilde{\mathcal{N}}_{L/M} \circ \langle\varphi\rangle_{L/M}(U) \cdot U_{\mathbb{X}(M/K)}, \quad (2.23)$$

for every $U \in U_{\tilde{\mathbb{X}}(L/K)}^\diamond$, where $\overline{U}$ denotes, as before, the coset $U \cdot U_{\mathbb{X}(L/K)}$ in $U_{\tilde{\mathbb{X}}(L/K)}^\diamond / U_{\mathbb{X}(L/K)}$.

Lemma 2.10. *For an infinite Galois sub-extension M/K of L/K such that the residue-class degree $[\kappa_M : \kappa_K] = d'$ and $K \subset M \subset K_{\varphi^{d'}}$ for some $d' \mid d$, the square*

$$\begin{array}{ccc} \text{Gal}(L/L_0) & \xrightarrow{\phi_{L/L_0}^{(\varphi^d)}} & U_{\tilde{\mathbb{X}}(L/L_0)}^\diamond / U_{\mathbb{X}(L/L_0)} \\ \text{res}_M \downarrow & & \downarrow \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} \\ \text{Gal}(M/M_0) & \xrightarrow{\phi_{M/M_0}^{(\varphi^{d'})}} & U_{\tilde{\mathbb{X}}(M/M_0)}^\diamond / U_{\mathbb{X}(M/M_0)}, \end{array} \quad (2.24)$$

where the right-vertical arrow is the Coleman norm map $\tilde{\mathcal{N}}_{L/M}^{\text{Coleman}}$ from L to M defined by eq.s (2.22) and (2.23), is commutative.

Proof. For $\sigma \in \text{Gal}(L/L_0)$, $\text{res}_M(\sigma) = \sigma|_M \in \text{Gal}(M/M_0)$, as $L_0 \cap M = L \cap K^{nr} \cap M = M \cap K^{nr} = M_0$. Now, for any $\sigma \in \text{Gal}(L/L_0)$, following the definition $\phi_{L/L_0}^{(\varphi^d)}(\sigma) = U_\sigma \cdot U_{\mathbb{X}(L/L_0)}$, where $U_\sigma \in U_{\mathbb{X}(L/L_0)}^\diamond$ satisfies the equation $U_\sigma^{1-\varphi^d} = \Pi_{\varphi^d; L/L_0}^{\sigma-1}$. Thus, to prove the commutativity of the square, it suffices to prove that

$$\tilde{\mathcal{N}}_{L/M}(U_\sigma^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}) \equiv U_{\sigma|_M} \pmod{U_{\mathbb{X}(M/M_0)}},$$

where $U_{\sigma|_M} \in U_{\mathbb{X}(M/M_0)}^\diamond$ satisfies the equation $U_{\sigma|_M}^{1-\varphi^{d'}} = \Pi_{\varphi^{d'}; M/M_0}^{\sigma|_M-1}$. Now, without loss of generality, in view of Remark 2.8, fix a *basic sequence* (cf. [8])

$$L_0 = E_0 \subset E_1 \subset \dots \subset E_i \subset \dots \subset L,$$

where

- (i) $L = \bigcup_{0 \leq i \in \mathbb{Z}} E_i$;
- (ii) E_i/L_0 is a Galois extension for every $0 \leq i \in \mathbb{Z}$;
- (iii) E_{i+1}/E_i is cyclic of prime degree $[E_{i+1} : E_i] = p = \text{char}(\kappa_{L_0})$ for each $1 \leq i \in \mathbb{Z}$;
- (iv) E_1/E_0 is cyclic of degree relatively prime to p .

Thus, each extension E_i/L_0 is finite and Galois for $0 \leq i \in \mathbb{Z}$. Now, note that

$$\tilde{\mathcal{N}}_{L/M}(U_\sigma^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)})^{1-\varphi^{d'}} = \tilde{\mathcal{N}}_{L/M}(U_\sigma)^{1-\varphi^d} = \tilde{\mathcal{N}}_{L/M}(U_\sigma^{1-\varphi^d}).$$

As $U_\sigma^{1-\varphi^d} = \Pi_{\varphi^d; L/L_0}^{\sigma-1}$, setting $U_\sigma = (u_{\tilde{E}_i})_{0 \leq i \in \mathbb{Z}}$, for $0 \leq i \in \mathbb{Z}$,

$$\begin{aligned} \tilde{\mathcal{N}}_{L/M}(U_\sigma^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)})^{1-\varphi^{d'}} &= \tilde{\mathcal{N}}_{L/M} \left(\Pi_{\varphi^d; L/L_0}^{\sigma-1} \right)_i \\ &= \tilde{N}_{E_i/E_i \cap M}(\pi_{E_i}^{\sigma-1})^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}} \\ &= N_{E_i/E_i \cap M}(\pi_{E_i}^{\sigma-1}) \\ &= \pi_{E_i \cap M}^{\sigma|_M-1}. \end{aligned}$$

Now, it follows that, $\tilde{\mathcal{N}}_{L/M} \circ \langle \varphi \rangle_{L/M}(U_\sigma)^{1-\varphi^{d'}} = \Pi_{\varphi^{d'}; M/M_0}^{\sigma|_M-1}$, which yields the congruence $\tilde{\mathcal{N}}_{L/M} \circ \langle \varphi \rangle_{L/M}(U_\sigma) \equiv U_{\sigma|_M} \pmod{U_{\mathbb{X}(M/M_0)}}$ completing the proof. $\square$

So, we have the following theorem.

Theorem 2.11. *For an infinite Galois sub-extension M/K of L/K such that the residue-class degree $[\kappa_M : \kappa_K] = d'$ and $K \subset M \subset K_{\varphi^{d'}}$ for some $d' \mid d$, the square*

$$\begin{array}{ccc} \text{Gal}(L/K) & \xrightarrow{\phi_{L/K}^{(\varphi)}} & K^\times / N_{L_0/K} L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)} \\ \text{res}_M \downarrow & & \downarrow (e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}}) \\ \text{Gal}(M/K) & \xrightarrow{\phi_{M/K}^{(\varphi)}} & K^\times / N_{M_0/K} M_0^\times \times U_{\mathbb{X}(M/K)}^\diamond / U_{\mathbb{X}(M/K)}, \end{array}$$

where the right-vertical arrow

$$K^\times / N_{L_0/K} L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)} \xrightarrow{(e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman})} K^\times / N_{M_0/K} M_0^\times \times U_{\mathbb{X}(M/K)}^\diamond / U_{\mathbb{X}(M/K)}$$

defined by

$$(e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman}) : (\bar{a}, \bar{U}) \mapsto (e_{L_0/M_0}^{CFT}(\bar{a}), \tilde{\mathcal{N}}_{L/M}^{Coleman}(\bar{U}))$$

for every $(\bar{a}, \bar{U}) \in K^\times / N_{L_0/K} L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)}$, is commutative. Here,

$$e_{L_0/M_0}^{CFT} : K^\times / N_{L_0/K} L_0^\times \rightarrow K^\times / N_{M_0/K} M_0^\times$$

is the natural inclusion defined via the existence theorem of local class field theory.

Proof. By the isomorphism defined by eq.s (2.1) and (2.2), for $\sigma \in \text{Gal}(L/K)$, there exists a unique $0 \leq m \in \mathbb{Z}$ such that $\sigma|_{L_0} = \varphi^m$ and $\varphi^{-m}\sigma \in \text{Gal}(L/L_0)$. Now, following the definition,

$$\phi_{L/K}^{(\varphi)}(\sigma) = \left(\pi_K^m N_{L_0/K} L_0^\times, \phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma) \right).$$

Thus,

$$\begin{aligned} (e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman}) \left(\pi_K^m N_{L_0/K} L_0^\times, \phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma) \right) &= \\ (e_{L_0/M_0}^{CFT}(\pi_K^m N_{L_0/K} L_0^\times), \tilde{\mathcal{N}}_{L/M}^{Coleman}(\phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma))) &= \\ \left(\pi_K^m N_{M_0/K} M_0^\times, \phi_{M/M_0}^{(\varphi^{d'})}(\varphi^{-m}\sigma|_M) \right) \end{aligned}$$

by Lemma 2.10. Note that, by the existence theorem of local class field theory,

$$e_{L_0/M_0}^{CFT}(\pi_K^m N_{L_0/K} L_0^\times) = \pi_K^m N_{M_0/K} M_0^\times = \pi_K^{m'} N_{M_0/K} M_0^\times,$$

where $0 \leq m' \in \mathbb{Z}$ is the unique integer satisfying $(\sigma|_M)|_{M_0} = \sigma|_{M_0} = \varphi^{m'}$ and $\varphi^{-m'}(\sigma|_M) \in \text{Gal}(M/M_0)$. Hence,

$$\begin{aligned} (e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman}) (\phi_{L/K}^{(\varphi)}(\sigma)) &= \left(\pi_K^{m'} N_{M_0/K} M_0^\times, \phi_{M/M_0}^{(\varphi^{d'})}(\varphi^{-m}\sigma|_M) \right) \\ &= \left(\pi_K^{m'} N_{M_0/K} M_0^\times, \phi_{M/M_0}^{(\varphi^{d'})}(\varphi^{-m'}(\sigma|_M)) \right) \\ &= \phi_{M/K}^{(\varphi)}(\text{res}_M(\sigma)) \end{aligned}$$

by Remark 2.2 part (i), which completes the proof. $\square$


Now, let F/K be a finite sub-extension of L/K . Thus, L/F is an infinite APF-Galois extension (cf. Lemma 3.3 of [8]). Fix a Lubin-Tate splitting φ_F over F . Now, assume that the residue-class degree $[\kappa_L : \kappa_F] = d'$, for some $d' \mid d$, and there exists the chain of field extensions

$$F \subset L \subset F_{(\varphi_F)^{d'}}.$$

Thus, there exists the generalized arrow

$$\phi_{L/F}^{(\varphi_F)} : \text{Gal}(L/F) \rightarrow F^\times / N_{L_0^{(F)}/F} L_0^{(F)\times} \times U_{\widetilde{\mathbb{X}}(L/F)}^\diamond / U_{\mathbb{X}(L/F)}$$

corresponding to the extension L/F . Here, $L_0^{(F)}$ is defined as usual by $L_0^{(F)} = L \cap F^{nr} = F_{d'}^{nr}$. Note that, there is the following diagram of field extensions.



Thus, $L/L_0^{(F)}$ and $L/L_0^{(K)}$ are infinite totally-ramified *APF*-Galois extensions, by Lemma 3.3 of [8], and satisfy $L_0^{(F)} \subset L \subset (L_0^{(F)})_{\varphi_F^{d'}}$ and $L_0^{(K)} \subset L \subset (L_0^{(K)})_{\varphi_K^d}$.

Remark 2.12. Note that $L_0^{(F)}$ is compatible with $(L_0^{(K)}, \varphi_{L_0^{(K)}})$, in the sense of [10] pp. 89, where $\varphi_{L_0^{(K)}} = \varphi_K^d$. Thus, $\varphi_{L_0^{(F)}} = \varphi_F^{d'} = \varphi_{L_0^{(K)}}^{f(L_0^{(F)}/L_0^{(K)})} = \varphi_K^d$, as $L_0^{(F)}/L_0^{(K)}$ is totally-ramified.

For the extension $L/L_0^{(F)}$, fix an ascending chain

$$L_0^{(F)} = F_0 \subset F_1 \subset \cdots \subset F_i \subset \cdots \subset L$$

satisfying $L = \bigcup_{0 \leq i \in \mathbb{Z}} F_i$ and $[F_{i+1} : F_i] < \infty$ for every $0 \leq i \in \mathbb{Z}$. Following [8], introduce the homomorphism

$$\Lambda_{F/K} : \widetilde{\mathbb{X}}(L/L_0^{(F)})^\times \rightarrow \widetilde{\mathbb{X}}(L/L_0^{(K)})^\times \quad (2.25)$$

by

$$\begin{aligned}
 \Lambda_{F/K} : (\alpha_{F_0} \xleftarrow{\widetilde{N}_{F_1/F_0}} \alpha_{F_1} \xleftarrow{\widetilde{N}_{F_2/F_1}} \cdots) \mapsto \\
 (\widetilde{N}_{L_0^{(F)}/L_0^{(K)}}(\alpha_{F_0}) \xleftarrow{\widetilde{N}_{L_0^{(F)}/L_0^{(K)}}} \alpha_{F_0} \xleftarrow{\widetilde{N}_{F_1/F}} \alpha_{F_1} \xleftarrow{\widetilde{N}_{F_2/F_1}} \cdots), \quad (2.26)
 \end{aligned}$$

for each $(\alpha_{F_i})_{0 \leq i \in \mathbb{Z}} \in \widetilde{\mathbb{X}}(L/L_0^{(F)})^\times$. This homomorphism induces a group homomorphism

$$\lambda_{F/K} : U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / U_{\mathbb{X}(L/L_0^{(F)})} \rightarrow U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / U_{\mathbb{X}(L/L_0^{(K)})} \quad (2.27)$$

defined by

$$\lambda_{F/K} : \overline{U} \mapsto \Lambda_{F/K}(U).U_{\mathbb{X}(L/L_0^{(K)})}, \quad (2.28)$$

for every $U \in U_{\mathbb{X}(L/L_0^{(F)})}^\diamond$, where $\overline{U}$ denotes the coset $U.U_{\mathbb{X}(L/L_0^{(F)})}$ in $U_{\mathbb{X}(L/L_0^{(F)})}^\diamond/U_{\mathbb{X}(L/L_0^{(F)})}$ (for details cf. [8]).

Lemma 2.13. *Let F/K be a finite sub-extension of L/K . Fix a Lubin-Tate splitting φ_F over F . Assume that the residue-class degree $[\kappa_L : \kappa_F] = d'$ and $F \subset L \subset F_{(\varphi_F)^{d'}}$ for some $d' \mid d$. Then the square*

$$\begin{array}{ccc} \text{Gal}(L/L_0^{(F)}) & \xrightarrow{\phi_{L/L_0^{(F)}}^{(\varphi_K^d)}} & U_{\mathbb{X}(L/L_0^{(F)})}^\diamond/U_{\mathbb{X}(L/L_0^{(F)})} \\ \text{inc.} \downarrow & & \downarrow \lambda_{F/K} \\ \text{Gal}(L/L_0^{(K)}) & \xrightarrow{\phi_{L/L_0^{(K)}}^{(\varphi_K^d)}} & U_{\mathbb{X}(L/L_0^{(K)})}^\diamond/U_{\mathbb{X}(L/L_0^{(K)})} \end{array} \quad (2.29)$$

where the right-vertical arrow

$$\lambda_{F/K} : U_{\mathbb{X}(L/L_0^{(F)})}^\diamond/U_{\mathbb{X}(L/L_0^{(F)})} \rightarrow U_{\mathbb{X}(L/L_0^{(K)})}^\diamond/U_{\mathbb{X}(L/L_0^{(K)})}$$

is defined by eq.s (2.27) and (2.28), is commutative.

Proof. Look at the proof of Theorem 5.12 of [8]. $\square$

So, we have the following theorem.

Theorem 2.14. *Let F/K be a finite sub-extension of L/K . Fix a Lubin-Tate splitting φ_F over F . Assume that the residue-class degree $[\kappa_L : \kappa_F] = d'$ and $F \subset L \subset F_{(\varphi_F)^{d'}}$ for some $d' \mid d$. Then the square*

$$\begin{array}{ccc} \text{Gal}(L/F) & \xrightarrow{\phi_{L/F}^{(\varphi_F)}} & F^\times/N_{L_0^{(F)}/F}L_0^{(F)\times} \times U_{\mathbb{X}(L/F)}^\diamond/U_{\mathbb{X}(L/F)} \\ \text{inc.} \downarrow & & \downarrow (N_{F/K}, \lambda_{F/K}) \\ \text{Gal}(L/K) & \xrightarrow{\phi_{L/K}^{(\varphi_K)}} & K^\times/N_{L_0^{(K)}/K}L_0^{(K)\times} \times U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)} \end{array} \quad (2.30)$$

where the right-vertical arrow

$$(N_{F/K}, \lambda_{F/K}) : F^\times/N_{L_0^{(F)}/F}L_0^{(F)\times} \times U_{\mathbb{X}(L/F)}^\diamond/U_{\mathbb{X}(L/F)} \rightarrow K^\times/N_{L_0^{(K)}/K}L_0^{(K)\times} \times U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)}$$

defined by

$$(N_{F/K}, \lambda_{F/K}) : (\overline{a}, \overline{U}) \mapsto \left(\overline{N_{F/K}(a)}, \lambda_{F/K}(\overline{U}) \right),$$

for every $(\overline{a}, \overline{U}) \in F^\times/N_{L_0^{(F)}/F}L_0^{(F)\times} \times U_{\mathbb{X}(L/F)}^\diamond/U_{\mathbb{X}(L/F)}$, is commutative.

Proof. Let $\sigma \in \text{Gal}(L/F)$. There exists $0 \leq m \in \mathbb{Z}$ such that $\sigma|_{L_0^{(F)}} = \varphi_F^m$ and $\varphi_F^{-m}\sigma \in \text{Gal}(L/L_0^{(F)})$. Now,

$$\phi_{L/F}^{(\varphi_F)}(\sigma) = \left(\pi_F^m \cdot N_{L_0^{(F)}/F} L_0^{(F)\times}, \phi_{L/L_0^{(F)}}^{(\varphi_K^d)}(\varphi_F^{-m} \sigma) \right)$$

and

$$(N_{F/K}, \lambda_{F/K})(\phi_{L/F}^{(\varphi_F)}(\sigma)) = \left(\pi_K^m \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}, \phi_{L/L_0^{(K)}}^{(\varphi_K^d)}(\varphi_F^{-m} \sigma) \right)$$

by the norm-compatibility of primes in the fixed Lubin-Tate labelling and by Lemma 2.13. Now, there exists $0 \leq m' \in \mathbb{Z}$ such that $\sigma|_{L_0^{(K)}} = \varphi_K^{m'}$ and $\varphi_K^{-m'}\sigma \in \text{Gal}(L/L_0^{(K)})$. By Remark 2.2 part (ii), it follows that $\varphi_F^m|_{L_0^{(K)}} = \varphi_K^{m'}$ and $\varphi_F^{-m}\sigma = \varphi_K^{-m'}\sigma$. By abelian local class field theory, $N_{F/K} : \pi_F^m N_{L_0^{(F)}/F} L_0^{(F)\times} \mapsto \pi_K^{m'} N_{L_0^{(K)}/K} L_0^{(K)\times} = \pi_K^m \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}$. Thus,

$$\begin{aligned} (N_{F/K}, \lambda_{F/K})(\phi_{L/F}^{(\varphi_F)}(\sigma)) &= \left(\pi_K^m \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}, \phi_{L/L_0^{(K)}}^{(\varphi_K^d)}(\varphi_F^{-m}\sigma) \right) \\ &= \left(\pi_K^{m'} \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}, \phi_{L/L_0^{(K)}}^{(\varphi_K^d)}(\varphi_K^{-m'}\sigma) \right) \\ &= \phi_{L/K}^{(\varphi_K)}(\sigma), \end{aligned}$$

which completes the proof. $\square$

Let L/K be any *APF*-Galois sub-extension of K_{φ^d}/K , where the residue-class degree is d . In case L/K is assumed to be a finite extension, the $\tilde{K}$ -coordinate of the generalized arrow $\phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \rightarrow K^\times/N_{L_0/K}L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\diamond/U_{\mathbb{X}(L/K)}$ is the Iwasawa-Neukirch map $\iota_{L/K}$ of L/K (for details on Iwasawa-Neukirch map $\iota_{L/K}$ of the Galois extension L/K , cf. Section 1 of [8]). More precisely, we have the following proposition.

Proposition 2.15. *Define a homomorphism*

$$\rho: K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/U_{\mathbb{X}(L/K)} \rightarrow K^\times/N_{L/K}L^\times \quad (2.31)$$

by

$$\rho : (\pi_K^m, (u_{\tilde{E}})) \mapsto \pi_K^m N_{L_0/K}(u_{\tilde{L}_0}) \pmod{N_{L/K} L^\times}, \quad (2.32)$$

for every $(\pi_K^m, (u_{\tilde{E}})) \in K^\times / N_{L_0/K} L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\circ / U_{\mathbb{X}(L/K)}$. Then the composite map

$$\begin{array}{c}
 \rho \circ \phi_{L/K}^{(\varphi)} = \iota_{L/K} \\
 \text{Gal}(L/K) \xrightarrow{\phi_{L/K}^{(\varphi)}} K^\times / N_{L_0/K} L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)} \xrightarrow{\rho} K^\times / N_{L/K} L^\times
 \end{array}
 \quad (2.33)$$

is the Iwasawa-Neukirch map $\iota_{L/K} : \text{Gal}(L/K) \rightarrow K^\times / N_{L/K} L^\times$ of L/K .

Proof. Let us briefly recall, following Section 1 of [8], the construction of the Iwasawa-Neukirch map

$$\iota_{L/K} : \text{Gal}(L/K) \rightarrow K^\times / N_{L/K} L^\times$$

for the Galois extension L/K . For each $\sigma \in \text{Gal}(L/K)$, choose $\sigma^* \in \text{Gal}(L^{nr}/K)$ in such a way that:

- (i) $\sigma^*|_L = \sigma$;
- (ii) $\sigma^*|_{K^{nr}} = \varphi^n$, for some $0 < n \in \mathbb{Z}$.

Let Σ_{σ^*} be the fixed-field $(L^{nr})^{\sigma^*}$ of $\sigma^* \in \text{Gal}(L^{nr}/K)$ in L^{nr} . It is well-known that $[\Sigma_{\sigma^*} : K] < \infty$. Now, the map $\iota_{L/K} : \text{Gal}(L/K) \rightarrow K^\times / N_{L/K} L^\times$ is defined by $\iota_{L/K}(\sigma) = N_{\Sigma_{\sigma^*}/K}(\pi_{\Sigma_{\sigma^*}}) \bmod N_{L/K} L^\times$, for $\sigma \in \text{Gal}(L/K)$, where $\pi_{\Sigma_{\sigma^*}}$ denotes any prime element of Σ_{σ^*} . Thus, for a finite *APF*-Galois extension L/K satisfying $[\kappa_L : \kappa_K] = d$ and $K \subset L \subset K_{\varphi^d}$, it suffices to prove that, for $\sigma \in \text{Gal}(L/K)$,

$$\rho \circ \phi_{L/K}^{(\varphi)}(\sigma) = \iota_{L/K}(\sigma) = N_{\Sigma_{\sigma^*}/K}(\pi_{\Sigma_{\sigma^*}}) \bmod N_{L/K} L^\times,$$

where $\pi_{\Sigma_{\sigma^*}}$ denotes any prime element of Σ_{σ^*} . Now, for $\sigma \in \text{Gal}(L/K)$, there exists $0 \leq m \in \mathbb{Z}$ such that $\sigma|_{L_0} = \varphi^m$ and $\tau = \varphi^{-m}\sigma \in \text{Gal}(L/L_0)$. Thus, $\sigma = \varphi^m \tau$.

Case 1: $m > 0$. In this case, it suffices to prove that

$$\pi_K^m N_{L_0/K} \left(\text{Pr}_{\tilde{L}_0}(\phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma)) \right) = N_{\Sigma_{\sigma^*}/K}(\pi_{\Sigma_{\sigma^*}}) \bmod N_{L/K} L^\times,$$

where $\pi_{\Sigma_{\sigma^*}}$ denotes any prime element of Σ_{σ^*} . To prove this equality, choose $\sigma^* \in \text{Gal}(L^{nr}/K)$ such that

- (i) $\sigma^*|_L = \sigma$;
- (ii) $\sigma^*|_{K^{nr}} = \varphi^m$.

In fact, let $\sigma^* = \varphi^m|_{L^{nr}} \tau^*$, where $\tau^* \in \text{Gal}(L^{nr}/L_0)$ is defined uniquely by the conditions $\tau^*|_L = \tau$ and $\tau^*|_{K^{nr}} = \text{id}_{K^{nr}}$, as $L^{nr} = LK^{nr}$. Note that, for $\Sigma = \Sigma_{\sigma^*}$, $\Sigma_0 = \Sigma \cap K^{nr}$ is a finite extension of degree $[\Sigma_0 : K] = m$ over K , as $\Sigma_0 = (K^{nr})^{\varphi^m}$, the fixed field of $\varphi^m \in \text{Gal}(K^{nr}/K)$ in K^{nr} . As L is fixed by φ^d , $T = L \cap K_\varphi$ is an unramified extension of degree d . Thus, the prime element π_T is a prime element of L and of L^{nr} . Now, choose a prime element π_Σ of Σ . It is well-known that $\Sigma^{nr} = L^{nr}$ (cf. Section 2 in Chapter 4 of [4]). Thus, π_Σ is a prime element of L^{nr} . So, there exists a unit $v \in L^{nr} \subset \tilde{L}$, such that $\pi_\Sigma = \pi_T v$. Note that, $\pi_\Sigma^{\sigma^*-1} = 1$ as Σ is fixed by σ^* . Thus, $(\pi_T v)^{\sigma^*-1} = 1$ and we get the equalities

$$\pi_T^{\sigma-1} = v^{1-\sigma^*} = v^{1-\varphi^m \tau^*} = v^{1-\tau^*} v^{(1-\varphi^m)\tau^*}.$$

Recall that (by Proposition 1.8 of Chapter IV of [4] or by 1.1 of [10]), $U_{\tilde{L}}$ is multiplicatively $(1 - \varphi^m)$ -divisible. So, there exists $w \in U_{\tilde{L}}$ such that $w^{1-\varphi^m} = v$. Hence,

$$\pi_T^{\sigma-1} = (w^{1-\tau^*} v^{\tau^*})^{1-\varphi^m},$$

as $\varphi^m \tau^* = \tau^* \varphi^m$. Now, choose $z \in U_{\tilde{L}}$ as $z = (w^{1-\tau^*} v^{\tau^*})^{1+\varphi+\dots+\varphi^{m-1}}$. Note that, this $z \in U_{\tilde{L}}$ satisfies $z^{1-\varphi} = \pi_T^{\sigma-1}$. Clearly,

$$\tilde{N}_{L/K}(z) = \tilde{N}_{L/K}(v)^{1+\varphi+\dots+\varphi^{m-1}}.$$

After this preliminary observations,

$$\begin{aligned} N_{\Sigma/K}(\pi_{\Sigma}) &= N_{\Sigma_0/K} \circ N_{\Sigma/\Sigma_0}(\pi_{\Sigma}) \\ &= \tilde{N}_{L/K}(\pi_{\Sigma})^{1+\varphi+\dots+\varphi^{m-1}} \\ &= \tilde{N}_{L/K}(\pi_T v)^{1+\sigma+\dots+\sigma^{m-1}} \\ &= \pi_K^m \tilde{N}_{L/K}(v)^{1+\varphi+\dots+\varphi^{m-1}} \\ &= \pi_K^m \tilde{N}_{L/K}(z), \end{aligned}$$

as π_T belongs to the fixed Lubin-Tate labelling. Thus, the image of σ under the Iwasawa-Neukirch map $\iota_{L/K}$ is

$$\iota_{L/K}(\sigma) = \pi_K^m \tilde{N}_{L/K}(z) \pmod{N_{L/K} L^{\times}}.$$

Now, let $y \in U_L$ such that

$$y^{1-\varphi^d} = \pi_T^{\varphi^{-m}\sigma-1} = \pi_T^{\sigma-1}.$$

Note that, $T = L \cap K_{\varphi}$. Then, setting $z = y^{1+\varphi+\dots+\varphi^{d-1}} \in U_L$,

$$z^{1-\varphi} = y^{1-\varphi^d} = \pi_T^{\sigma-1}.$$

Thus,

$$N_{L/K}(y) = \tilde{N}_{L/K}(y)^{1+\varphi+\dots+\varphi^{d-1}} = \tilde{N}_{L/K}(z),$$

which shows that

$$\begin{aligned} \iota_{L/K}(\sigma) &= \pi_K^m \tilde{N}_{L/K}(z) \pmod{N_{L/K} L^{\times}} \\ &= \pi_K^m N_{L/K}(y) \pmod{N_{L/K} L^{\times}} \\ &= \pi_K^m N_{L_0/K} \left(\tilde{N}_{L/K}(y) \right) \pmod{N_{L/K} L^{\times}} \\ &= \pi_K^m N_{L_0/K} \left(\text{Pr}_{\tilde{L}_0}(\phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma)) \right) \\ &= \rho \circ \phi_{L/K}^{(\varphi)}(\sigma), \end{aligned}$$

completing the proof.

Case 2: $m = 0$. In this case, $\sigma \in \text{Gal}(L/L_0)$. Consider $\varphi^d \sigma \in \text{Gal}(L^{nr}/K)$. Then by the previous Case 1,

$$\iota_{L/K}(\varphi^d \sigma) = \rho \circ \phi_{L/K}^{(\varphi)}(\varphi^d \sigma),$$

where $\iota_{L/K}(\varphi^d \sigma) = \iota_{L/K}(\sigma)$. Now, by Theorem 2.4,

$$\phi_{L/K}^{(\varphi)}(\varphi^d \sigma) = \phi_{L/K}^{(\varphi)}(\varphi^d) \phi_{L/K}^{(\varphi)}(\sigma) \varphi^d = \left(\pi_K^d \cdot N_{L_0/K} L_0^{\times}, \phi_{L/L_0}^{(\varphi^d)}(\sigma) \right),$$

where the last equality follows from the fact that $K \subset L \subset K_{\varphi^d}$. Thus,

$$\rho \circ \phi_{L/K}^{(\varphi)}(\sigma) = \rho \circ \phi_{L/K}^{(\varphi)}(\varphi^d \sigma) = \pi_K^d N_{L_0/K} \left(\text{Pr}_{\tilde{L}_0}(\phi_{L/L_0}^{(\varphi^d)}(\sigma)) \right) \pmod{N_{L/K} L^\times},$$

which proves that

$$\iota_{L/K}(\sigma) = \rho \circ \phi_{L/K}^{(\varphi)}(\sigma),$$

completing the proof. $\square$

Now, we shall generalize the definition of the extended Hazewinkel map $H_{L/K}^{(\varphi)} : U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/K} \rightarrow \text{Gal}(L/K)$ of Fesenko (cf. [1, 2, 3] and [8]) initially defined for totally-ramified *APF*-Galois sub-extensions L/K of K_φ/K to infinite *APF*-Galois sub-extensions L/K of K_{φ^d}/K , where $[\kappa_L : \kappa_K] = d$.

In order to do so, we first have to assume that the local field K satisfies the condition

$$\mu_p(K^{sep}) = \{\alpha \in K^{sep} : \alpha^p = 1\} \subset K, \quad (2.34)$$

where $p = \text{char}(\kappa_K)$. For details on the assumption (2.34) on K , we refer the reader to [1, 2, 3].

Let L/K be an infinite *APF*-Galois extension with residue-class degree $[\kappa_L : \kappa_K] = d$ and $K \subset L \subset K_{\varphi^d}$. As usual, let $L_0 = L \cap K^{nr}$. Define the generalized arrow

$$\mathbf{H}_{L/K}^{(\varphi)} : K^\times / N_{L_0/K} L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0} \rightarrow \text{Gal}(L/K) \quad (2.35)$$

for the extension L/K by

$$\mathbf{H}_{L/K}^{(\varphi)}((\pi_K^m N_{L_0/K} L_0^\times, U.Y_{L/L_0})) = \varphi^m |_L H_{L/L_0}^{(\varphi^d)}(U.Y_{L/L_0}), \quad (2.36)$$

for every $m \in \mathbb{Z}$ and $U \in U_{\tilde{\mathbb{X}}(L/K)}^\diamond$, where $H_{L/L_0}^{(\varphi^d)} : U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0} \rightarrow \text{Gal}(L/L_0)$ is the extended Hazewinkel map of Fesenko for the extension L/L_0 . For the definition and basic properties of the group Y_{L/L_0} , we refer the reader to [3] and [8].

The following lemma is clear.

Lemma 2.16. *Suppose that the local field K satisfies the condition given in eq. (2.34). Let L/K be an infinite *APF*-Galois sub-extension of K_{φ^d}/K , where $d = [\kappa_L : \kappa_K]$. Then the generalized arrow*

$$\mathbf{H}_{L/K}^{(\varphi)} : K^\times / N_{L_0/K} L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0} \rightarrow \text{Gal}(L/K) \quad (2.37)$$

for the extension L/K is a bijection.

Proof. The proof follows from the isomorphism $\text{Gal}(L/K) \simeq \text{Gal}(L_0/K) \times \text{Gal}(L/L_0)$ combined with the bijectivity of $H_{L/L_0}^{(\varphi^d)} : U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0} \rightarrow \text{Gal}(L/L_0)$ (cf. Lemma 5.22 of [8]) and abelian local class field theory for the extension L_0/K . $\square$

Now, consider the composition of arrows

$$\begin{array}{ccc}
\mathrm{Gal}(L/K) & \xrightarrow{\Phi_{L/K}^{(\varphi)}} & K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\circ/U_{\mathbb{X}(L/K)} \\
& \searrow \Phi_{L/K}^{(\varphi)} & \downarrow (\mathrm{id}_{K^\times/N_{L_0/K}L_0^\times}, c_{L/L_0}) \\
& & K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\circ/Y_{L/L_0}
\end{array} \quad (2.38)$$

where $c_{L/L_0} : U_{\mathbb{X}(L/K)}^\circ/U_{\mathbb{X}(L/K)} \rightarrow U_{\mathbb{X}(L/K)}^\circ/Y_{L/L_0}$ is the canonical map defined via the inclusion $U_{\mathbb{X}(L/K)} \subseteq Y_{L/L_0}$. Recall that (cf. eq. 5.35 of [8]), the composition $c_{L/L_0} \circ \phi_{L/L_0}^{(\varphi^d)} = \Phi_{L/L_0}^{(\varphi^d)} : \mathrm{Gal}(L/L_0) \rightarrow U_{\mathbb{X}(L/K)}^\circ/Y_{L/L_0}$ is the reciprocity map of Fesenko for the extension L/L_0 . Now, let $\sigma \in \mathrm{Gal}(L/K)$. Let $0 \leq m \in \mathbb{Z}$ such that $\sigma|_{L_0} = \varphi^m|_{L_0}$ and $\varphi^{-m}\sigma \in \mathrm{Gal}(L/L_0)$. Then following the definition,

$$\begin{aligned}
H_{L/K}^{(\varphi)} \circ \Phi_{L/K}^{(\varphi)}(\sigma) &= H_{L/K}^{(\varphi)} \left(\pi_K^m \cdot N_{L_0/K}L_0^\times, c_{L/L_0} \circ \phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma) \right) \\
&= H_{L/K}^{(\varphi)} \left(\pi_K^m \cdot N_{L_0/K}L_0^\times, \Phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma) \right) \\
&= \varphi^m|_L H_{L/L_0}^{(\varphi^d)} \left(\Phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma) \right) \\
&= \varphi^m|_L (\varphi^{-m}\sigma) \\
&= \sigma,
\end{aligned}$$

by Lemma 5.23 of [8]. For $0 \leq m \in \mathbb{Z}$ and $U \in U_{\mathbb{X}(L/K)}^\circ$, let $(\pi_K^m \cdot N_{L_0/K}L_0^\times, U \cdot Y_{L/L_0}) \in K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\circ/Y_{L/L_0}$. Now, again following the definition,

$$\begin{aligned}
\Phi_{L/K}^{(\varphi)} \circ H_{L/K}^{(\varphi)}((\pi_K^m \cdot N_{L_0/K}L_0^\times, U \cdot Y_{L/L_0})) &= \Phi_{L/K}^{(\varphi)} \left(\varphi^m|_L H_{L/L_0}^{(\varphi^d)}(U \cdot Y_{L/L_0}) \right) \\
&= (\mathrm{id}_{K^\times/N_{L_0/K}L_0^\times}, c_{L/L_0}) \circ \Phi_{L/K}^{(\varphi)} \left(\varphi^m|_L H_{L/L_0}^{(\varphi^d)}(U \cdot Y_{L/L_0}) \right) \\
&= (\mathrm{id}_{K^\times/N_{L_0/K}L_0^\times}, c_{L/L_0}) \left(\pi_K^m \cdot N_{L_0/K}L_0^\times, \phi_{L/L_0}^{(\varphi^d)}(H_{L/L_0}^{(\varphi^d)}(U \cdot Y_{L/L_0})) \right) \\
&= \left(\pi_K^m \cdot N_{L_0/K}L_0^\times, \Phi_{L/L_0}^{(\varphi^d)}(H_{L/L_0}^{(\varphi^d)}(U \cdot Y_{L/L_0})) \right) \\
&= (\pi_K^m \cdot N_{L_0/K}L_0^\times, U \cdot Y_{L/L_0}),
\end{aligned}$$

by Lemma 5.23 of [8]. Thus, these computations yield

$$H_{L/K}^{(\varphi)} \circ \Phi_{L/K}^{(\varphi)} = \mathrm{id}_{\mathrm{Gal}(L/K)} \quad (2.39)$$

and

$$\Phi_{L/K}^{(\varphi)} \circ H_{L/K}^{(\varphi)} = \mathrm{id}_{K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\circ/Y_{L/L_0}}. \quad (2.40)$$

Note that, there is a natural continuous action of $\mathrm{Gal}(L/K)$ on the topological group $K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\circ/Y_{L/L_0}$ defined by abelian local class field theory on the first component and by eq. (5.7) and Lemma 5.20 of [8] on the second component as

$$(\overline{a}, \overline{U})^\sigma = \left(\overline{a}^{\varphi^m}, \overline{U}^{\varphi^{-m}\sigma} \right) = \left(\overline{a}, \overline{U}^{\varphi^{-m}\sigma} \right), \quad (2.41)$$

for every $\sigma \in \text{Gal}(L/K)$, where $\sigma|_{L_0} = \varphi^m$ for some $0 \leq m \in \mathbb{Z}$, and for every $a \in K^\times$ with $\bar{a} = a.N_{L_0/K}L_0^\times$ and $U \in U_{\mathbb{X}(L/K)}^\diamond$ with $\bar{U} = U.Y_{L/L_0}$. We shall always view $K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0}$ as a topological $\text{Gal}(L/K)$ -module in this text.

So, we have the following theorem, which follows from Theorem 2.4, Lemma 2.16 and eq.s (2.39) and (2.40) combined with the fact that $U_{\mathbb{X}(L/K)}$ is a topological $\text{Gal}(L/L_0)$ -submodule of Y_{L/L_0} .

Theorem 2.17. *Suppose that the local field K satisfies the condition given in eq. (2.34). Let L/K be an infinite APF-Galois sub-extension of K_{φ^d}/K , where $d = [\kappa_L : \kappa_K]$. The mapping*

$$\Phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \rightarrow K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0}$$

defined for the extension L/K is a bijection with the inverse

$$H_{L/K}^{(\varphi)} : K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0} \rightarrow \text{Gal}(L/K).$$

For every $\sigma, \tau \in \text{Gal}(L/K)$,

$$\Phi_{L/K}^{(\varphi)}(\sigma\tau) = \Phi_{L/K}^{(\varphi)}(\sigma)\Phi_{L/K}^{(\varphi)}(\tau)^\sigma \quad (2.42)$$

co-cycle condition is satisfied.

By Corollary 2.5, Theorem 2.17 has the following consequence.

Corollary 2.18. *Define a law of composition $*$ on $K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0}$ by*

$$(\bar{a}, \bar{U}) * (\bar{b}, \bar{V}) = (\bar{a}, \bar{U}).(\bar{b}, \bar{V})^{(\Phi_{L/K}^{(\varphi)})^{-1}((\bar{a}, \bar{U}))} \quad (2.43)$$

for every $\bar{a} = a.N_{L_0/K}L_0^\times, \bar{b} = b.N_{L_0/K}L_0^\times \in K^\times/N_{L_0/K}L_0^\times$ with $a, b \in K^\times$ and $\bar{U} = U.Y_{L/L_0}, \bar{V} = V.Y_{L/L_0} \in U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0}$ with $U, V \in U_{\mathbb{X}(L/K)}^\diamond$. Then $K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0}$ is a topological group under $*$, and the map $\Phi_{L/K}^{(\varphi)}$ induces an isomorphism of topological groups

$$\Phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \xrightarrow{\sim} K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0}, \quad (2.44)$$

where the topological group structure on $K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0}$ is defined with respect to the binary operation $*$ defined by eq. (2.43).

Definition 2.19. Let K be a local field satisfying the condition given in eq. (2.34). Let L/K be an infinite APF-Galois sub-extension of K_{φ^d}/K , where $d = [\kappa_L : \kappa_K]$. The mapping

$$\Phi_{L/K}^{(\varphi)} : \text{Gal}(L/K) \rightarrow K^\times/N_{L_0/K}L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond/Y_{L/L_0},$$

defined in Theorem 2.17, is called the *generalized Fesenko reciprocity map* for the extension L/K .

For each $0 \leq i \in \mathbb{R}$, we have previously introduced the higher unit groups $\left(U_{\tilde{\mathbb{X}}(L/K)}^\diamond\right)^i$ of the field $\tilde{\mathbb{X}}(L/K)$ as in eq. (2.16). For each $0 \leq n \in \mathbb{Z}$, as in eq. 5.42 of [8], let

$$Q_{L/L_0}^n = c_{L/L_0} \left(\left(U_{\tilde{\mathbb{X}}(L/K)}^\diamond \right)^n U_{\tilde{\mathbb{X}}(L/K)} / U_{\tilde{\mathbb{X}}(L/K)} \cap \text{im}(\phi_{L/L_0}^{(\varphi^d)}) \right), \quad (2.45)$$

which is a subgroup of $\left(U_{\tilde{\mathbb{X}}(L/K)}^\diamond\right)^n Y_{L/L_0} / Y_{L/L_0}$. Now, ramification theorem, stated in Theorem 2.7, can be reformulated for the generalized reciprocity map $\Phi_{L/K}^{(\varphi)}$ corresponding to the extension L/K as follows.

Theorem 2.20 (Ramification theorem). *Let K be a local field satisfying the condition given in eq. (2.34). For $0 \leq u \in \mathbb{R}$, let $\text{Gal}(L/K)_u$ denote the u^{th} higher ramification subgroup in the lower numbering of the Galois group $\text{Gal}(L/K)$ corresponding to the infinite APF-Galois sub-extension L/K of K_{φ^d}/K with residue-class degree $[\kappa_L : \kappa_K] = d$. Then, for $0 \leq n \in \mathbb{Z}$, there exists the inclusion*

$$\begin{aligned} \Phi_{L/K}^{(\varphi)} \left(\text{Gal}(L/K)_{\psi_{L/K} \circ \varphi_{L/L_0}(n)} - \text{Gal}(L/K)_{\psi_{L/K} \circ \varphi_{L/L_0}(n+1)} \right) \subseteq \\ \left\langle 1_{K^\times / N_{L_0/K} L_0^\times} \right\rangle \times \left(\left(U_{\tilde{\mathbb{X}}(L/K)}^\diamond \right)^n Y_{L/L_0} / Y_{L/L_0} - Q_{L/L_0}^{n+1} \right). \end{aligned}$$

Proof. Following the general observation made in the first paragraph of the proof of Theorem 2.7, for $0 \leq u \in \mathbb{R}$ and for $\tau \in \text{Gal}(L/K)_u = \text{Gal}(L/L_0)^{\varphi_{L/K}(u)}$,

$$\phi_{L/K}^{(\varphi)}(\tau) = \left(1_{K^\times / N_{L_0/K} L_0^\times}, \phi_{L/L_0}^{(\varphi')}(\tau) \right),$$

where $\varphi' = \varphi^d$. Thus, following the definition,

$$\begin{aligned} \Phi_{L/K}^{(\varphi)}(\tau) &= \left(1_{K^\times / N_{L_0/K} L_0^\times}, c_{L/L_0} \circ \phi_{L/L_0}^{(\varphi')}(\tau) \right) \\ &= \left(1_{K^\times / N_{L_0/K} L_0^\times}, \Phi_{L/L_0}^{(\varphi')}(\tau) \right). \end{aligned}$$

Now, to prove the theorem, let $u = \psi_{L/K} \circ \varphi_{L/L_0}(n)$ and $u' = \psi_{L/K} \circ \varphi_{L/L_0}(n+1)$. Then, for any $\tau \in \text{Gal}(L/K)_u - \text{Gal}(L/K)_{u'}$, it follows from the ramification theorem (cf. Theorem 5.27 in [8]) that the second coordinate of $\Phi_{L/K}^{(\varphi)}(\tau)$ satisfies

$$\Phi_{L/L_0}^{(\varphi')}(\tau) \in \left(U_{\tilde{\mathbb{X}}(L/K)}^\diamond \right)^n Y_{L/L_0} / Y_{L/L_0} - Q_{L/L_0}^{n+1},$$

since

$$\text{Gal}(L/K)_u = \text{Gal}(L/L_0)^{\varphi_{L/K}(u)} = \text{Gal}(L/L_0)^{\varphi_{L/L_0}(n)} = \text{Gal}(L/L_0)_n$$

and likewise

$$\text{Gal}(L/K)_{u'} = \text{Gal}(L/L_0)^{\varphi_{L/K}(u')} = \text{Gal}(L/L_0)^{\varphi_{L/L_0}(n+1)} = \text{Gal}(L/L_0)_{n+1},$$

which completes the proof. $\square$

Let K be a local field satisfying the condition given in eq. (2.34). Let M/K be an infinite Galois sub-extension of L/K . Thus, by Lemma 3.3 of [8], M is an *APF*-Galois extension over K . We further assume that, the residue-class degree $[\kappa_M : \kappa_K] = d'$ and $K \subset M \subset K_{\varphi^{d'}}$ for some $d' \mid d$. Let

$$\Phi_{M/K}^{(\varphi)} : \text{Gal}(M/K) \rightarrow K^\times / N_{M_0/K} M_0^\times \times U_{\tilde{\mathbb{X}}(M/K)}^\diamond / Y_{M/M_0}$$

be the corresponding generalized Fesenko reciprocity map defined for the extension M/K . Here, M_0 is defined as usual by $M_0 = M \cap K^{nr} = K_{d'}^{nr}$. Now, fix a basic sequence

$$L_o = E_o \subset E_1 \subset \cdots \subset E_i \subset \cdots \subset L$$

for the extension L/L_o . Now, following the notation of [3] and [8], introduce for each $1 \leq i \in \mathbb{Z}$, an element σ_i in $\text{Gal}(\tilde{L}/\tilde{K})$ such that $\langle \sigma |_{E_i} \rangle = \text{Gal}(E_i/E_{i-1})$. Further, for each $1 \leq k, i \in \mathbb{Z}$, introduce the map $h_k^{(L/L_o)} : \prod_{1 \leq i \leq k} U_{\tilde{E}_k}^{\sigma_i-1} \rightarrow \left(\prod_{1 \leq i \leq k+1} U_{\tilde{E}_{k+1}}^{\sigma_i-1} \right) / U_{\tilde{E}_{k+1}}^{\sigma_{k+1}-1}$, the map $g_k^{(L/L_o)} : \prod_{1 \leq i \leq k} U_{\tilde{E}_k}^{\sigma_i-1} \rightarrow \prod_{1 \leq i \leq k+1} U_{\tilde{E}_{k+1}}^{\sigma_i-1}$ and the map $f_i^{(L/L_o)} : U_{\tilde{E}_i}^{\sigma_i-1} \rightarrow U_{\tilde{\mathbb{X}}(L/E_i)} \xrightarrow{\Lambda_{E_i/E_o}} U_{\tilde{\mathbb{X}}(L/K)}$ following [3] and [8]. Now, fix the sequence

$$M_o = E_o \cap M \subseteq E_1 \cap M \subseteq \cdots \subseteq E_i \cap M \subseteq \cdots \subseteq M$$

for the extension M/M_o . Define, for each $1 \leq k \in \mathbb{Z}$, a homomorphism

$$h_k^{(M/M_o)} : \prod_{1 \leq i \leq k} U_{\widetilde{E_k \cap M}}^{\sigma_i|_{\tilde{M}}-1} \rightarrow \left(\prod_{1 \leq i \leq k+1} U_{\widetilde{E_{k+1} \cap M}}^{\sigma_i|_{\tilde{M}}-1} \right) / U_{\widetilde{E_{k+1} \cap M}}^{\sigma_{k+1}|_{\tilde{M}}-1}$$

that satisfies

$$\left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \tilde{N}_{E_{k+1}/E_{k+1} \cap M} \right) \circ h_k^{(L/L_o)} = h_k^{(M/M_o)} \circ \left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \tilde{N}_{E_k/E_k \cap M} \right)$$

and any map

$$g_k^{(M/M_o)} : \prod_{1 \leq i \leq k} U_{\widetilde{E_k \cap M}}^{\sigma_i|_{\tilde{M}}-1} \rightarrow \prod_{1 \leq i \leq k+1} U_{\widetilde{E_{k+1} \cap M}}^{\sigma_i|_{\tilde{M}}-1}$$

that satisfies

$$\left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \tilde{N}_{E_{k+1}/E_{k+1} \cap M} \right) \circ g_k^{(L/L_o)} = g_k^{(M/M_o)} \circ \left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \tilde{N}_{E_k/E_k \cap M} \right)$$

following the same lines of [3] and [8].

Now, for each $1 \leq i \in \mathbb{Z}$, introduce the map

$$f_i^{(M/M_o)} : U_{\widetilde{E_i \cap M}}^{\sigma_i|_{\tilde{M}}-1} \rightarrow U_{\tilde{\mathbb{X}}(M/K)}$$

by

$$f_i^{(M/M_o)}(w) = \tilde{N}_{L/M} \left(f_i^{(L/L_o)}(v) \right),$$

where $v \in U_{\widetilde{E}_i}^{\sigma_i-1}$ is any element satisfying $\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_i/E_i \cap M}(v) = w \in U_{\widetilde{E}_i \cap M}^{\sigma_i|_{\widetilde{M}}-1}$. Note that, if $v' \in U_{\widetilde{E}_i}^{\sigma_i-1}$ such that $\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_i/E_i \cap M}(v') = w$, then $\widetilde{N}_{L/M}(f_i^{(L/L_o)}(v)) = \widetilde{N}_{L/M}(f_i^{(L/L_o)}(v'))$.

In fact, there exists $u \in \ker \left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_i/E_i \cap M} \right)$ such that $v' = vu$. Thus, we have to verify that $\widetilde{N}_{L/M}(f_i^{(L/L_o)}(v)) = \widetilde{N}_{L/M}(f_i^{(L/L_o)}(vu))$. That is, for each $1 \leq j \in \mathbb{Z}$, we have to check that

$$\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_j/E_j \cap M} \left(\text{Pr}_{\widetilde{E}_j}(f_i^{(L/L_o)}(v)) \right) = \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_j/E_j \cap M} \left(\text{Pr}_{\widetilde{E}_j}(f_i^{(L/L_o)}(vu)) \right).$$

Now, for $j > i$, it follows that

$$\begin{aligned} & \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_j/E_j \cap M} \left(\text{Pr}_{\widetilde{E}_j}(f_i^{(L/L_o)}(v)) \right) = \\ & \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_j/E_j \cap M} \left(g_{j-1}^{(L/L_o)} \circ \dots \circ g_i^{(L/L_o)}(v) \right) = \\ & g_{j-1}^{(M/M_o)} \circ \dots \circ g_i^{(M/M_o)} \left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_i/E_i \cap M}(v) \right) = \\ & g_{j-1}^{(M/M_o)} \circ \dots \circ g_i^{(M/M_o)} \left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_i/E_i \cap M}(vu) \right) = \\ & \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_j/E_j \cap M} \left(g_{j-1}^{(L/L_o)} \circ \dots \circ g_i^{(L/L_o)}(vu) \right) = \\ & \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_j/E_j \cap M} \left(\text{Pr}_{\widetilde{E}_j}(f_i^{(L/L_o)}(vu)) \right). \end{aligned}$$

Thus, the map

$$f_i^{(M/M_o)} : U_{\widetilde{E}_i \cap M}^{\sigma_i|_{\widetilde{M}}-1} \rightarrow U_{\widetilde{\mathbb{X}}(M/K)}$$

is well-defined. Moreover, for $j > i$,

$$\text{Pr}_{\widetilde{E}_j \cap M} \circ f_i^{(M/M_o)} = \left(g_{j-1}^{(M/M_o)} \circ \dots \circ g_i^{(M/M_o)} \right) \big|_{U_{\widetilde{E}_i \cap M}^{\sigma_i|_{\widetilde{M}}-1}}.$$

In fact, for $w \in U_{\widetilde{E}_i \cap M}^{\sigma_i|_{\widetilde{M}}-1}$, there exists $v \in U_{\widetilde{E}_i}^{\sigma_i-1}$ such that $\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_i/E_i \cap M}(v) = w$, and $f_i^{(M/M_o)}(w) = \widetilde{N}_{L/M}(f_i^{(L/L_o)}(v))$. That is, the following square

$$\begin{array}{ccc} U_{\widetilde{E}_i}^{\sigma_i-1} & \xrightarrow{f_i^{(L/L_o)}} & U_{\widetilde{\mathbb{X}}(L/K)} \\ \downarrow \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \widetilde{N}_{E_i/E_i \cap M} & & \downarrow \widetilde{N}_{L/M} \\ U_{\widetilde{E}_i \cap M}^{\sigma_i|_{\widetilde{M}}-1} & \xrightarrow{f_i^{(M/M_o)}} & U_{\widetilde{\mathbb{X}}(M/K)} \end{array} \quad (2.46)$$

is commutative. Thus,

$$\begin{aligned}
\Pr_{\widetilde{E_j \cap M}} \circ f_i^{(M/M_o)}(w) &= \Pr_{\widetilde{E_j \cap M}} \circ \tilde{\mathcal{N}}_{L/M} \left(f_i^{(L/L_o)}(v) \right) \\
&= \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \tilde{\mathcal{N}}_{E_j/E_j \cap M} \left(\Pr_{\tilde{E_j}} \circ f_i^{(L/L_o)}(v) \right) \\
&= \prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \tilde{\mathcal{N}}_{E_j/E_j \cap M} \left((g_{j-1}^{(L/L_o)} \circ \dots \circ g_i^{(L/L_o)})(v) \right) \\
&= \left(g_{j-1}^{(M/M_o)} \circ \dots \circ g_i^{(M/M_o)} \right) \left(\prod_{0 \leq \ell \leq f(L/M)-1} (\varphi^{d'})^\ell \tilde{\mathcal{N}}_{E_i/E_i \cap M}(v) \right),
\end{aligned}$$

which is the desired equality.

Now, we shall modify Lemma 5.28 of [8] and show that the norm map $\tilde{\mathcal{N}}_{L/M} : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times$ introduced by eq.s (2.17) and (2.18) satisfies the following lemma.

Lemma 2.21. *The norm map $\tilde{\mathcal{N}}_{L/M} : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times$ introduced by eq.s (2.17) and (2.18) further satisfies*

- (i) $\tilde{\mathcal{N}}_{L/M} \left(Z_{L/L_o} \left(\{E_i, f_i^{(L/L_o)}\} \right) \right) \subseteq Z_{M/M_o} \left(\{E_i \cap M, f_i^{(M/M_o)}\} \right);$
- (ii) $\tilde{\mathcal{N}}_{L/M} \circ \langle \varphi \rangle_{L/M} (Y_{L/L_o}) \subseteq Y_{M/M_o}.$

Proof. (i) Recall that, $\tilde{\mathcal{N}}_{L/M} : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times$ is a continuous mapping. Now, for any choice of $z^{(i)} \in \text{im}(f_i^{(L/L_o)})$, the continuity of the multiplicative arrow $\tilde{\mathcal{N}}_{L/M} : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times$ yields

$$\tilde{\mathcal{N}}_{L/M} \left(\prod_i z^{(i)} \right) = \prod_i \tilde{\mathcal{N}}_{L/M}(z^{(i)}),$$

where $\tilde{\mathcal{N}}_{L/M}(z^{(i)}) \in \text{im}(f_i^{(M/M_o)})$ by the commutative square (2.46).

- (ii) For $y \in Y_{L/L_o}$, as $y^{1-\varphi^d} \in Z_{L/L_o}$, it follows that $\tilde{\mathcal{N}}_{L/M}(y^{1-\varphi^d}) \in Z_{M/M_o}$ by part (i). Now, note that

$$\tilde{\mathcal{N}}_{L/M}(y^{1-\varphi^d}) = \tilde{\mathcal{N}}_{L/M}(y)^{1-\varphi^d} = \left(\tilde{\mathcal{N}}_{L/M}(y)^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}} \right)^{1-\varphi^{d'}}.$$

Therefore,

$$\tilde{\mathcal{N}}_{L/M}(y)^{1+\varphi^{d'}+\dots+\varphi^{d'(f(L/M)-1)}} = \tilde{\mathcal{N}}_{L/M} \circ \langle \varphi \rangle_{L/M} (y) \in Y_{M/M_o}$$

as desired.

The proof is now complete. $\square$

Thus, by part (ii) of Lemma 2.21, the homomorphism $\tilde{\mathcal{N}}_{L/M} \circ \langle \varphi \rangle_{L/M} : \tilde{\mathbb{X}}(L/K)^\times \rightarrow \tilde{\mathbb{X}}(M/K)^\times$ induces a group homomorphism, which will again be called the *Coleman norm map from L to M*,

$$\tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} : U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_o} \rightarrow U_{\tilde{\mathbb{X}}(M/K)}^\diamond / Y_{M/M_o} \quad (2.47)$$

and defined by

$$\tilde{\mathcal{N}}_{L/M}^{\text{Coleman}}(\overline{U}) = \tilde{\mathcal{N}}_{L/M} \circ \langle \varphi \rangle_{L/M}(U) \cdot Y_{M/M_0}, \quad (2.48)$$

for every $U \in U_{\mathbb{X}(L/K)}^\diamond$, where $\overline{U}$ denotes, as usual, the coset $U \cdot Y_{L/L_0}$ in $U_{\mathbb{X}(L/K)}^\diamond / Y_{L/L_0}$.

The following lemma is the finer version of Lemma 2.10.

Lemma 2.22. *Let K be a local field satisfying the condition given in eq. (2.34). For an infinite Galois sub-extension M/K of L/K such that the residue-class degree $[\kappa_M : \kappa_K] = d'$ and $K \subset M \subset K_{\varphi^{d'}}$ for some $d' \mid d$, the square*

$$\begin{array}{ccc} \text{Gal}(L/L_0) & \xrightarrow{\Phi_{L/L_0}^{(\varphi^d)}} & U_{\mathbb{X}(L/L_0)}^\diamond / Y_{L/L_0} \\ \text{res}_M \downarrow & & \downarrow \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} \\ \text{Gal}(M/M_0) & \xrightarrow{\Phi_{M/M_0}^{(\varphi^{d'})}} & U_{\mathbb{X}(M/M_0)}^\diamond / Y_{M/M_0}, \end{array} \quad (2.49)$$

where the right-vertical arrow is the Coleman norm map $\tilde{\mathcal{N}}_{L/M}^{\text{Coleman}}$ from L to M defined by eq.s (2.47) and (2.48), is commutative.

Proof. It suffices to prove that the square

$$\begin{array}{ccc} U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)} & \xrightarrow{c_{L/L_0}} & U_{\mathbb{X}(L/K)}^\diamond / Y_{L/L_0} \\ \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} \downarrow & & \downarrow \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} \\ U_{\mathbb{X}(M/K)}^\diamond / U_{\mathbb{X}(M/K)} & \xrightarrow{c_{M/M_0}} & U_{\mathbb{X}(M/K)}^\diamond / Y_{M/M_0} \end{array}$$

is commutative, which is obvious. Then pasting this square with the square eq. (2.24) as

$$\begin{array}{ccccc} \text{Gal}(L/K) & \xrightarrow{\phi_{L/L_0}^{(\varphi^d)}} & U_{\mathbb{X}(L/K)}^\diamond / U_{\mathbb{X}(L/K)} & \xrightarrow{c_{L/L_0}} & U_{\mathbb{X}(L/K)}^\diamond / Y_{L/K} \\ \text{res}_M \downarrow & & \downarrow \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} & & \downarrow \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}} \\ \text{Gal}(M/K) & \xrightarrow{\phi_{M/M_0}^{(\varphi^{d'})}} & U_{\mathbb{X}(M/K)}^\diamond / U_{\mathbb{X}(M/K)} & \xrightarrow{c_{M/M_0}} & U_{\mathbb{X}(M/K)}^\diamond / Y_{M/K} \end{array}$$

the commutativity of the square eq. (2.49) follows. $\square$

Thus, we have the following theorem, which is the finer version of Theorem 2.11.

Theorem 2.23. *Let K be a local field satisfying the condition given in eq. (2.34). For an infinite Galois sub-extension M/K of L/K such that the residue-class degree $[\kappa_M : \kappa_K] = d'$ and $K \subset M \subset K_{\varphi^{d'}}$ for some $d' \mid d$, the square*

$$\begin{array}{ccc} \text{Gal}(L/K) & \xrightarrow{\Phi_{L/K}^{(\varphi)}} & K^\times / N_{L_0/K} L_0^\times \times U_{\mathbb{X}(L/K)}^\diamond / Y_{L/L_0} \\ \text{res}_M \downarrow & & \downarrow (e_{L_0/M_0}^{\text{CFT}}, \tilde{\mathcal{N}}_{L/M}^{\text{Coleman}}) \\ \text{Gal}(M/K) & \xrightarrow{\Phi_{M/K}^{(\varphi)}} & K^\times / N_{M_0/K} M_0^\times \times U_{\mathbb{X}(M/K)}^\diamond / Y_{M/M_0}, \end{array}$$

where the right-vertical arrow

$$K^\times / N_{L_0/K} L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0} \xrightarrow{(e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman})} K^\times / N_{M_0/K} M_0^\times \times U_{\tilde{\mathbb{X}}(M/K)}^\diamond / Y_{M/M_0}$$

defined by

$$(e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman}) : (\bar{a}, \bar{U}) \mapsto (e_{L_0/M_0}^{CFT}(\bar{a}), \tilde{\mathcal{N}}_{L/M}^{Coleman}(\bar{U}))$$

for every $(\bar{a}, \bar{U}) \in K^\times / N_{L_0/K} L_0^\times \times U_{\tilde{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0}$, is commutative. Here,

$$e_{L_0/M_0}^{CFT} : K^\times / N_{L_0/K} L_0^\times \rightarrow K^\times / N_{M_0/K} M_0^\times$$

is the natural inclusion defined via the existence theorem of local class field theory.

Proof. By the isomorphism defined by eq.s (2.1) and (2.2), for $\sigma \in \text{Gal}(L/K)$, there exists a unique $0 \leq m \in \mathbb{Z}$ such that $\sigma|_{L_0} = \varphi^m$ and $\varphi^{-m}\sigma \in \text{Gal}(L/L_0)$. Now, following the definition,

$$\Phi_{L/K}^{(\varphi)}(\sigma) = \left(\pi_K^m N_{L_0/K} L_0^\times, \Phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma) \right).$$

Thus,

$$\begin{aligned} & \left(e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman} \right) \left(\pi_K^m N_{L_0/K} L_0^\times, \Phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma) \right) = \\ & \left(e_{L_0/M_0}^{CFT}(\pi_K^m N_{L_0/K} L_0^\times), \tilde{\mathcal{N}}_{L/M}^{Coleman}(\Phi_{L/L_0}^{(\varphi^d)}(\varphi^{-m}\sigma)) \right) = \\ & \left(\pi_K^m N_{M_0/K} M_0^\times, \Phi_{M/M_0}^{(\varphi^{d'})}(\varphi^{-m}\sigma|_M) \right) \end{aligned}$$

by Lemma 2.22. Note that, by the existence theorem of local class field theory,

$$e_{L_0/M_0}^{CFT}(\pi_K^m N_{L_0/K} L_0^\times) = \pi_K^m N_{M_0/K} M_0^\times = \pi_K^{m'} N_{M_0/K} M_0^\times,$$

where $0 \leq m' \in \mathbb{Z}$ is the unique integer satisfying $(\sigma|_M)|_{M_0} = \sigma|_{M_0} = \varphi^{m'}$ and $\varphi^{-m'}(\sigma|_M) \in \text{Gal}(M/M_0)$. Hence,

$$\begin{aligned} & \left(e_{L_0/M_0}^{CFT}, \tilde{\mathcal{N}}_{L/M}^{Coleman} \right) (\Phi_{L/K}^{(\varphi)}(\sigma)) = \left(\pi_K^{m'} N_{M_0/K} M_0^\times, \Phi_{M/M_0}^{(\varphi^{d'})}(\varphi^{-m}\sigma|_M) \right) \\ & = \left(\pi_K^{m'} N_{M_0/K} M_0^\times, \Phi_{M/M_0}^{(\varphi^{d'})}(\varphi^{-m'}(\sigma|_M)) \right) \\ & = \Phi_{M/K}^{(\varphi)}(\text{res}_M(\sigma)) \end{aligned}$$

by Remark 2.2 part (i), which completes the proof. $\square$

Let K be a local field satisfying the condition given in eq. (2.34). Now, let F/K be a finite sub-extension of L/K . Thus, L/F is an infinite *APF*-Galois extension (cf. Lemma 3.3 of [8]), where F satisfies (2.34). Fix a Lubin-Tate splitting φ_F over F . Now, assume that the residue-class degree $[\kappa_L : \kappa_F] = d'$, for some $d' \mid d$, and there exists the chain of field extensions

$$F \subset L \subset F_{(\varphi_F)^{d'}}.$$

Thus, there exists the generalized Fesenko reciprocity map

$$\Phi_{L/F}^{(\varphi_F)} : \text{Gal}(L/F) \rightarrow F^\times / N_{L_0^{(F)}/F} L_0^{(F)\times} \times U_{\mathbb{X}(L/F)}^\diamond / Y_{L/L_0^{(F)}}$$

corresponding to the extension L/F . Here, $L_0^{(F)}$ is defined as usual by $L_0^{(F)} = L \cap F^{nr} = F_{d'}^{nr}$ (and recall that, $L_0^{(K)} = L \cap K^{nr} = K_d^{nr}$).

Now, fix a basic sequence

$$L_0^{(K)} = E_o \subset E_1 \subset \cdots \subset E_i \subset \cdots \subset L$$

for the extension $L/L_0^{(K)}$. Following the notation of [3] and [8], introduce for each $1 \leq i \in \mathbb{Z}$, an element σ_i in $\text{Gal}(\tilde{L}/\tilde{K})$ such that $\langle \sigma |_{E_i} \rangle = \text{Gal}(E_i/E_{i-1})$. Now, fix the sequence

$$L_0^{(F)} = E_o L_0^{(F)} \subseteq E_1 L_0^{(F)} \subseteq \cdots \subseteq E_i L_0^{(F)} \subseteq \cdots \subseteq L L_0^{(F)} = L$$

for the extension $L/L_0^{(F)}$ following eq. (5.55) of [8]. Now, for $1 \leq i \in \mathbb{Z}$, introduce the elements σ_i^* in $\text{Gal}(\tilde{L}/\tilde{F})$ that satisfies

$$\langle \sigma_i^* |_{E_i L_0^{(F)}} \rangle = \text{Gal}(E_i L_0^{(F)} / E_{i-1} L_0^{(F)})$$

as follows :

(i) in case $i > i_o$, then define $\sigma_i^* = \sigma_i$;

(ii) in case $i \leq i_o$, then define

$$\sigma_i^* = \begin{cases} \sigma_i, & E_{i-1} L_0^{(F)} \subset E_i L_0^{(F)}; \\ \text{id}_{E_i L_0^{(F)}}, & E_{i-1} L_0^{(F)} = E_i L_0^{(F)}; \end{cases}$$

where i_o is defined as in the paragaph of eq. (5.55) of [8]. It is then clear that, for each $1 \leq i \in \mathbb{Z}$, the elements σ_i^* of $\text{Gal}(\tilde{L}/\tilde{F})$ satisfies

$$\langle \sigma_i^* |_{E_i L_0^{(F)}} \rangle = \text{Gal}(E_i L_0^{(F)} / E_{i-1} L_0^{(F)}),$$

and for almost all i , $\sigma_i^* = \sigma_i$. Further, for each $1 \leq k, i \in \mathbb{Z}$, introduce the map $h_k^{(L/L_0^{(F)})} : \prod_{1 \leq i \leq k} \widetilde{U_{E_k L_0^{(F)}}^{\sigma_i^* - 1}} \rightarrow \left(\prod_{1 \leq i \leq k+1} \widetilde{U_{E_{k+1} L_0^{(F)}}^{\sigma_i^* - 1}} \right) / \widetilde{U_{E_{k+1} L_0^{(F)}}^{\sigma_{k+1}^* - 1}}$, the

map $g_k^{(L/L_0^{(F)})} : \prod_{1 \leq i \leq k} \widetilde{U_{E_k L_0^{(F)}}^{\sigma_i^* - 1}} \rightarrow \prod_{1 \leq i \leq k+1} \widetilde{U_{E_{k+1} L_0^{(F)}}^{\sigma_i^* - 1}}$ and the map $f_i^{(L/L_0^{(F)})} : \widetilde{U_{E_i L_0^{(F)}}^{\sigma_i^* - 1}} \rightarrow \widetilde{U_{\mathbb{X}(L/E_i L_0^{(F)})}^{\sigma_i^* - 1}} \xrightarrow{\Lambda_{E_i L_0^{(F)}/L_0^{(F)}}} \widetilde{U_{\mathbb{X}(L/F)}^{\sigma_i^* - 1}}$ following [3] and [8]. Define, for each $1 \leq k \in \mathbb{Z}$, a homomorphism

$$h_k^{(L/L_0^{(K)})} : \prod_{1 \leq i \leq k} \widetilde{U_{\tilde{E}_k}^{\sigma_i - 1}} \rightarrow \left(\prod_{1 \leq i \leq k+1} \widetilde{U_{\tilde{E}_{k+1}}^{\sigma_i - 1}} \right) / \widetilde{U_{\tilde{E}_{k+1}}^{\sigma_{k+1} - 1}}$$

that satisfies

$$\tilde{N}_{E_{k+1} L_0^{(F)}/E_{k+1}}^* \circ h_k^{(L/L_0^{(F)})} = h_k^{(L/L_0^{(K)})} \circ \tilde{N}_{E_k L_0^{(F)}/E_k}$$

and any map

$$g_k^{(L/L_0^{(K)})} : \prod_{1 \leq i \leq k} U_{\tilde{E}_k}^{\sigma_i - 1} \rightarrow \prod_{1 \leq i \leq k+1} U_{\tilde{E}_{k+1}}^{\sigma_i - 1}$$

that satisfies

$$\tilde{N}_{E_{k+1}L_0^{(F)}/E_{k+1}} \circ g_k^{(L/L_0^{(F)})} = g_k^{(L/L_0^{(K)})} \circ \tilde{N}_{E_kL_0^{(F)}/E_k}$$

following the same lines of [3] and [8].

Now, for each $1 \leq i \in \mathbb{Z}$, introduce the map

$$f_i^{(L/L_0^{(K)})} : U_{\tilde{E}_i}^{\sigma_i - 1} \rightarrow U_{\tilde{\mathbb{X}}(L/K)}$$

by

$$f_i^{(L/L_0^{(K)})}(w) = \Lambda_{F/K} \left(f_i^{(L/L_0^{(F)})}(v) \right),$$

where $v \in U_{E_iL_0^{(F)}}^{\sigma_i^* - 1}$ is any element satisfying $\tilde{N}_{E_iL_0^{(F)}/E_i}(v) = w \in U_{\tilde{E}_i}^{\sigma_i - 1}$. Note

that, if $v' \in U_{E_iL_0^{(F)}}^{\sigma_i^* - 1}$ such that $\tilde{N}_{E_iL_0^{(F)}/E_i}(v') = w$, then $\Lambda_{F/K} \left(f_i^{(L/L_0^{(F)})}(v) \right) = \Lambda_{F/K} \left(f_i^{(L/L_0^{(F)})}(v') \right)$.

In fact, there exists $u \in \ker \left(\tilde{N}_{E_iL_0^{(F)}/E_i} \right)$ such that $v' = vu$. Thus, we have to verify that $\Lambda_{F/K} \left(f_i^{(L/L_0^{(F)})}(v) \right) = \Lambda_{F/K} \left(f_i^{(L/L_0^{(F)})}(vu) \right)$. That is, for each $1 \leq j \in \mathbb{Z}$, we have to check that

$$\Pr_{\tilde{E}_j} \left(\Lambda_{F/K}(f_i^{(L/L_0^{(F)})}(v)) \right) = \Pr_{\tilde{E}_j} \left(\Lambda_{F/K}(f_i^{(L/L_0^{(F)})}(vu)) \right). \quad (2.50)$$

Now, for $j > i$, it follows that

$$\begin{aligned} \Pr_{\tilde{E}_j} \left(\Lambda_{F/K}(f_i^{(L/L_0^{(F)})}(v)) \right) &= \tilde{N}_{E_jL_0^{(F)}/E_j} \left(\Pr_{\tilde{E}_jL_0^{(F)}} \left(\Lambda_{F/K}(f_i^{(L/L_0^{(F)})}(v)) \right) \right) \\ &= \tilde{N}_{E_jL_0^{(F)}/E_j} \left(\Pr_{\tilde{E}_jL_0^{(F)}} \left(f_i^{(L/L_0^{(F)})}(v) \right) \right) \\ &= \tilde{N}_{E_jL_0^{(F)}/E_j} \left(g_{j-1}^{(L/L_0^{(F)})} \circ \dots \circ g_i^{(L/L_0^{(F)})}(v) \right) \\ &= g_{j-1}^{(L/L_0^{(K)})} \circ \dots \circ g_i^{(L/L_0^{(K)})} \left(\tilde{N}_{E_iL_0^{(F)}/E_i}(v) \right), \end{aligned}$$

by the properties of the mappings $g_k^{(L/L_0^{(F)})}$ and $g_k^{(L/L_0^{(K)})}$. Thus, eq. (2.50) follows, as $\tilde{N}_{E_iL_0^{(F)}/E_i}(v) = \tilde{N}_{E_iL_0^{(F)}/E_i}(vu)$. Therefore, the map

$$f_i^{(L/L_0^{(K)})} : U_{\tilde{E}_i}^{\sigma_i - 1} \rightarrow U_{\tilde{\mathbb{X}}(L/K)}$$

is well-defined. Moreover, for $j > i$,

$$\Pr_{\tilde{E}_j} \circ f_i^{(L/L_0^{(K)})} = \left(g_{j-1}^{(L/L_0^{(K)})} \circ \dots \circ g_i^{(L/L_0^{(K)})} \right) |_{U_{\tilde{E}_i}^{\sigma_i - 1}}.$$

In fact, for $w \in U_{\tilde{E}_i}^{\sigma_i-1}$, there exists $v \in \widetilde{U_{E_i L_0^{(F)}}^{\sigma_i^*-1}}$ such that $\tilde{N}_{E_i L_0^{(F)}/E_i}(v) = w$, and $f_i^{(L/L_0^{(K)})}(w) = \Lambda_{F/K} \left(f_i^{(L/L_0^{(F)})}(v) \right)$. That is, the following square

$$\begin{array}{ccc} \widetilde{U_{E_i L_0^{(F)}}^{\sigma_i^*-1}} & \xrightarrow{f_i^{(L/L_0^{(F)})}} & U_{\tilde{\mathbb{X}}(L/F)} \\ \tilde{N}_{E_i L_0^{(F)}/E_i} \downarrow & & \downarrow \Lambda_{F/K} \\ U_{\tilde{E}_i}^{\sigma_i-1} & \xrightarrow{f_i^{(L/L_0^{(K)})}} & U_{\tilde{\mathbb{X}}(L/K)} \end{array} \quad (2.51)$$

is commutative. Thus,

$$\begin{aligned} \Pr_{\tilde{E}_j} \circ f_i^{(L/L_0^{(K)})}(w) &= \Pr_{\tilde{E}_j} \circ \Lambda_{F/K} \left(f_i^{(L/L_0^{(F)})}(v) \right) \\ &= \tilde{N}_{E_j L_0^{(F)}/E_j} \left(\Pr_{\widetilde{U_{E_j L_0^{(F)}}}} \circ f_i^{(L/L_0^{(F)})}(v) \right) \\ &= \tilde{N}_{E_j L_0^{(F)}/E_j} \left((g_{j-1}^{(L/L_0^{(F)})} \circ \dots \circ g_i^{(L/L_0^{(F)})})(v) \right) \\ &= \left(g_{j-1}^{(L/L_0^{(K)})} \circ \dots \circ g_i^{(L/L_0^{(K)})} \right) \left(\tilde{N}_{E_i L_0^{(F)}/E_i}(v) \right), \end{aligned}$$

by the properties of the mappings $g_k^{(L/L_0^{(F)})}$ and $g_k^{(L/L_0^{(K)})}$, which is the desired equality.

Now, we shall modify Lemma 5.30 of [8] and show that the homomorphism $\Lambda_{F/K} : \tilde{\mathbb{X}}(L/L_0^{(F)})^\times \rightarrow \tilde{\mathbb{X}}(L/L_0^{(K)})^\times$ introduced by eq.s (2.25) and (2.26) satisfies the following lemma.

Lemma 2.24. *The continuous homomorphism $\Lambda_{F/K} : \tilde{\mathbb{X}}(L/L_0^{(F)})^\times \rightarrow \tilde{\mathbb{X}}(L/L_0^{(K)})^\times$ introduced by eq.s (2.25) and (2.26) further satisfies*

- (i) $\Lambda_{F/K} \left(Z_{L/L_0^{(F)}}(\{K_i F, f_i^{(L/L_0^{(F)})}\}) \right) \subseteq Z_{L/L_0^{(K)}}(\{K_i, f_i^{(L/L_0^{(K)})}\})$;
- (ii) $\Lambda_{F/K}(Y_{L/L_0^{(F)}}) \subseteq Y_{L/L_0^{(K)}}.$

Proof. (i) For any choice of $z^{(i)} \in Z_i^{(L/L_0^{(F)})}$, the continuity of the multiplicative arrow $\Lambda_{F/K} : \tilde{\mathbb{X}}(L/L_0^{(F)})^\times \rightarrow \tilde{\mathbb{X}}(L/L_0^{(K)})^\times$ yields

$$\Lambda_{F/K} \left(\prod_i z^{(i)} \right) = \prod_i \Lambda_{F/K}(z^{(i)}),$$

where $\Lambda_{F/K}(z^{(i)}) \in Z_i^{(L/L_0^{(K)})}$ by the commutative square (2.51).

- (ii) Now let $y \in Y_{L/L_0^{(F)}}$. Then, $y^{1-\varphi_F^{d'}} \in Z_{L/L_0^{(F)}}(\{K_i F, f_i^{(L/L_0^{(F)})}\})$. Thus, $\Lambda_{F/K}(y^{1-\varphi_F^{d'}}) = \Lambda_{F/K}(y)^{1-\varphi_F^{d'}} \in Z_{L/L_0^{(K)}}(\{K_i, f_i^{(L/L_0^{(K)})}\})$ by part (i). Now the result follows, as $\varphi_F^{d'} = \varphi_K^d$ by Remark 2.12. □

Thus, the homomorphism $\Lambda_{F/K} : \widetilde{\mathbb{X}}(L/L_0^{(F)})^\times \rightarrow \widetilde{\mathbb{X}}(L/L_0^{(K)})^\times$ of eq. (2.25) defined by eq. (2.26) induces a group homomorphism,

$$\lambda_{F/K} : U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / Y_{L/L_0^{(F)}} \rightarrow U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / Y_{L/L_0^{(K)}} \quad (2.52)$$

and defined by

$$\lambda_{F/K}(\overline{U}) = \Lambda_{F/K}(U) \cdot Y_{L/L_0^{(K)}}, \quad (2.53)$$

for every $U \in U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond$, where $\overline{U}$ denotes, as usual, the coset $U \cdot Y_{L/L_0^{(F)}}$ in $U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / Y_{L/L_0^{(F)}}$.

Let

$$\Phi_{L/F}^{(\varphi_F)} : \text{Gal}(L/F) \rightarrow F^\times / N_{L_0^{(F)}/F} L_0^{(F)\times} \times U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / Y_{L/L_0^{(F)}}$$

be the corresponding generalized Fesenko reciprocity map defined for the extension L/F , where $Y_{L/L_0^{(F)}} = Y_{L/L_0^{(F)}} \left(\{K_i F, f_i^{(L/L_0^{(F)})}\} \right)$.

The following lemma is the finer version of Lemma 2.13.

Lemma 2.25. *Let K be a local field satisfying the condition given in eq. (2.34). Let F/K be a finite sub-extension of L/K . Fix a Lubin-Tate splitting φ_F over F . Assume that the residue-class degree $[\kappa_L : \kappa_F] = d'$ and $F \subset L \subset F_{(\varphi_F)^{d'}}$ for some $d' \mid d$. Then the square*

$$\begin{array}{ccc} \text{Gal}(L/L_0^{(F)}) & \xrightarrow{\Phi_{L/L_0^{(F)}}^{(\varphi_K^d)}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / Y_{L/L_0^{(F)}} \\ \text{inc.} \downarrow & & \downarrow \lambda_{F/K} \\ \text{Gal}(L/L_0^{(K)}) & \xrightarrow{\Phi_{L/L_0^{(K)}}^{(\varphi_K^d)}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / Y_{L/L_0^{(K)}}, \end{array} \quad (2.54)$$

where the right-vertical arrow

$$\lambda_{F/K} : U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / Y_{L/L_0^{(F)}} \rightarrow U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / Y_{L/L_0^{(K)}}$$

is defined by eq.s (2.52) and (2.53), is commutative.

Proof. It suffices to prove that the square

$$\begin{array}{ccc} U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})} & \xrightarrow{\text{can.}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / Y_{L/L_0^{(F)}} \\ \lambda_{F/K} \downarrow & & \downarrow \lambda_{F/K} \\ U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})} & \xrightarrow{\text{can.}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / Y_{L/L_0^{(K)}} \end{array}$$

is commutative, which is obvious. Then pasting this square with the square eq. (2.29) as

$$\begin{array}{ccccc} \text{Gal}(L/L_0^{(F)}) & \xrightarrow{\Phi_{L/L_0^{(F)}}^{(\varphi_K^d)}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})} & \xrightarrow{\text{can.}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(F)})}^\diamond / Y_{L/L_0^{(F)}} \\ \text{inc.} \downarrow & & \downarrow \lambda_{L/F} & & \downarrow \lambda_{L/F} \\ \text{Gal}(L/L_0^{(K)}) & \xrightarrow{\Phi_{L/L_0^{(K)}}^{(\varphi_K^d)}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})} & \xrightarrow{\text{can.}} & U_{\widetilde{\mathbb{X}}(L/L_0^{(K)})}^\diamond / Y_{L/L_0^{(K)}} \end{array}$$

the commutativity of the square eq. (2.54) follows. $\square$

Thus, we have the following theorem, which is the finer version of Theorem 2.14.

Theorem 2.26. *Let K be a local field satisfying the condition given in eq. (2.34). Let F/K be a finite sub-extension of L/K . Fix a Lubin-Tate splitting φ_F over F . Assume that the residue-class degree $[\kappa_L : \kappa_F] = d'$ and $F \subset L \subset F_{(\varphi_F)^{d'}}$ for some $d' \mid d$. Then the square*

$$\begin{array}{ccc} \text{Gal}(L/F) & \xrightarrow{\Phi_{L/F}^{(\varphi_F)}} & F^\times / N_{L_0^{(F)}/F} L_0^{(F)\times} \times U_{\mathbb{X}(L/F)}^\diamond / Y_{L/L_0^{(F)}} \\ \text{inc.} \downarrow & & \downarrow (N_{F/K}, \lambda_{F/K}) \\ \text{Gal}(L/K) & \xrightarrow{\Phi_{L/K}^{(\varphi_K)}} & K^\times / N_{L_0^{(K)}/K} L_0^{(K)\times} \times U_{\mathbb{X}(L/K)}^\diamond / Y_{L/L_0^{(K)}}, \end{array} \quad (2.55)$$

where the right-vertical arrow

$$(N_{F/K}, \lambda_{F/K}) : F^\times / N_{L_0^{(F)}/F} L_0^{(F)\times} \times U_{\mathbb{X}(L/F)}^\diamond / Y_{L/L_0^{(F)}} \rightarrow K^\times / N_{L_0^{(K)}/K} L_0^{(K)\times} \times U_{\mathbb{X}(L/K)}^\diamond / Y_{L/L_0^{(K)}}$$

defined by

$$(N_{F/K}, \lambda_{F/K}) : (\bar{a}, \bar{U}) \mapsto \left(\overline{N_{F/K}(a)}, \lambda_{F/K}(\bar{U}) \right),$$

for every $(\bar{a}, \bar{U}) \in F^\times / N_{L_0^{(F)}/F} L_0^{(F)\times} \times U_{\mathbb{X}(L/F)}^\diamond / Y_{L/L_0^{(F)}}$, is commutative.

Proof. Let $\sigma \in \text{Gal}(L/F)$. There exists $0 \leq m \in \mathbb{Z}$ such that $\sigma|_{L_0^{(F)}} = \varphi_F^m$ and $\varphi_F^{-m}\sigma \in \text{Gal}(L/L_0^{(F)})$. Now,

$$\Phi_{L/F}^{(\varphi_F)}(\sigma) = \left(\pi_F^m \cdot N_{L_0^{(F)}/F} L_0^{(F)\times}, \Phi_{L/L_0^{(F)}}^{(\varphi_F^d)}(\varphi_F^{-m}\sigma) \right)$$

and

$$(N_{F/K}, \lambda_{F/K})(\Phi_{L/F}^{(\varphi_F)}(\sigma)) = \left(\pi_K^m \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}, \Phi_{L/L_0^{(K)}}^{(\varphi_K^d)}(\varphi_F^{-m}\sigma) \right)$$

by the norm-compatibility of primes in the fixed Lubin-Tate labelling and by Lemma 2.25. Now, there exists $0 \leq m' \in \mathbb{Z}$ such that $\sigma|_{L_0^{(K)}} = \varphi_K^{m'}$ and $\varphi_K^{-m'}\sigma \in \text{Gal}(L/L_0^{(K)})$. By Remark 2.2 part (ii), it follows that $\varphi_F^m|_{L_0^{(K)}} = \varphi_K^{m'}$ and $\varphi_F^{-m}\sigma = \varphi_K^{-m'}\sigma$. By abelian local class field theory, $N_{F/K} : \pi_F^m N_{L_0^{(F)}/F} L_0^{(F)\times} \mapsto \pi_K^{m'} N_{L_0^{(K)}/K} L_0^{(K)\times} = \pi_K^m \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}$. Thus,

$$\begin{aligned} (N_{F/K}, \lambda_{F/K})(\Phi_{L/F}^{(\varphi_F)}(\sigma)) &= \left(\pi_K^m \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}, \Phi_{L/L_0^{(K)}}^{(\varphi_K^d)}(\varphi_F^{-m}\sigma) \right) \\ &= \left(\pi_K^{m'} \cdot N_{L_0^{(K)}/K} L_0^{(K)\times}, \Phi_{L/L_0^{(K)}}^{(\varphi_K^d)}(\varphi_K^{-m'}\sigma) \right) \\ &= \Phi_{L/K}^{(\varphi_K)}(\sigma), \end{aligned}$$

which completes the proof. $\square$

Finally, the inverse $\mathbf{H}_{L/K}^{(\varphi)} = (\Phi_{L/K}^{(\varphi)})^{-1}$ of the generalized Fesenko reciprocity map $\Phi_{L/K}^{(\varphi)}$ defined for the extension L/K is the generalization of the Hazewinkel map for infinite APF -Galois sub-extensions L/K of K_{φ^d}/K satisfying $[\kappa_L : \kappa_K] = d$ and under the assumption that the local field K satisfies the condition given by eq. (2.34). More precisely, we have the following proposition.

Proposition 2.27. *The following square*

$$\begin{array}{ccc} K^\times / N_{L_0/K} L_0^\times \times U_{\widehat{\mathbb{X}}(L/K)}^\diamond / Y_{L/L_0} & \xrightarrow{\mathbf{H}_{L/K}^{(\varphi)}} & \text{Gal}(L/K) \\ \downarrow (id_{K^\times / N_{L_0/K} L_0^\times}, Pr_{\widehat{K}}) & & \downarrow \text{mod } \text{Gal}(L/K)' \\ K^\times / N_{L_0/K} L_0^\times \times U_{L_0}/N_{L/L_0} U_L & \xrightarrow{h_{L/K}} & \text{Gal}(L/K)^{ab} \end{array}$$

is commutative, where

$$h_{L/K} : K^\times / N_{L_0/K} L_0^\times \times U_{L_0}/N_{L/L_0} U_L \rightarrow \text{Gal}(L/K)^{ab}$$

is the Hazewinkel map of L/K .

References

- [1] I. B. Fesenko, *Local reciprocity cycles*, Invitation to Higher Local Fields (Ed. I. B. Fesenko, M. Kurihara), Geometry & Topology Monographs **3**, Warwick, 2000, 293-298
- [2] I. B. Fesenko, *Noncommutative local reciprocity maps*, Class Field Theory - Its Centenary and Prospect (Ed. K. Miyake), Advanced Studies in Pure Math. **30**, 2001, 63-78
- [3] I. B. Fesenko, *On the image of noncommutative local reciprocity map*, Homology, Homotopy and Appl. **7**, 2005, 53-62.
- [4] I. B. Fesenko, S. V. Vostokov, *Local Fields and Their Extensions: A Constructive Approach*, AMS Translations of Mathematical Monographs **121**, AMS, Providence, Rhode Island, 1993
- [5] J.-M. Fontaine, J.-P. Wintenberger, *Le “corps des normes” de certaines extensions algébriques de corps locaux*, C. R. Acad. Sci. Paris Sér. A Math. **288**, 1979, 367-370
- [6] J.-M. Fontaine, J.-P. Wintenberger, *Extensions algébriques et corps des normes des extensions APF des corps locaux*, C. R. Acad. Sci. Paris Sér. A Math. **288**, 1979, 441-444
- [7] A. Gurevich, *Ph.D. Thesis*, Humboldt Univ., Berlin, 1997
- [8] K. I. Ikeda and E. Serbest, *Fesenko reciprocity map*, submitted to Algebra i Analiz (2007).
- [9] K. I. Ikeda and E. Serbest, *Non-abelian local class field theory*, preprint (2007).

- [10] H. Koch and E. de Shalit, *Metabelian local class field theory*, J. reine angew. Math. **478**, 1996, 85-106
- [11] F. Laubie, *Une théorie du corps de classes local non abélien*, Compositio Math. **143**, 2007, 339-362.
- [12] J.-P. Wintenberger, *Le corps des normes de certaines extensions infinies de corps locaux; applications*, Ann. sci. Éc. Norm. Sup. **46**, 1983, 59-89

DEPARTMENT OF MATHEMATICS, ISTANBUL BILGI UNIVERSITY, KURTULUŞ
 DERESİ CAD. NO. 47, DOLAPDERE, 34440 BEYOĞLU, ISTANBUL, TURKEY
 E-mail : ilhan@bilgi.edu.tr
 GÜMÜŞ PALA MAHALLESİ, GÜMÜŞ SOK., ÖZ AKSU SİTESİ, C-2/39, 34160
 AVCILAR, ISTANBUL, TURKEY
 E-mail : erols73@yahoo.com